

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.

# APPLICATION UNDER UNITED STATES PATENT LAWS

Invention: USE OF PENETRATION ENHANCERS AND BARRIER DISRUPTION AGENTS TO ENHANCE THE TRANSCUTANEOUS IMMUNE RESPONSE INDUCED BY ADP-RIBOSYLATING EXOTOXIN

Inventor(s): Gregory M. GLENN  
Carl R. ALVING

Pillsbury Madison & Sutro LLP  
Intellectual Property Group  
1100 New York Avenue, N.W.  
Ninth Floor, East Tower  
Washington, D.C. 20005-3918  
Attorneys  
Telephone: (202) 861-3000

This is a:

- ☐ Provisional Application
- ☒ Regular Utility Application
- ☐ Continuing Application
- ☐ PCT National Phase Application
- ☐ Design Application
- ☐ Reissue Application
- ☐ Plant Application
- ☐ Substitute Specification  
Sub. Spec. filed  
in App. No. \_\_\_\_/
- ☐ Marked Up Specification re  
Sub. Spec. filed  
in App. No. \_\_\_\_/

## SPECIFICATION

USE OF PENETRATION ENHANCERS AND BARRIER DISRUPTION AGENTS  
TO ENHANCE THE TRANSCUTANEOUS IMMUNE RESPONSE INDUCED BY  
ADP-RIBOSYLATING EXOTOXIN

~~5~~ ~~1~~ ~~2~~ ~~3~~ ~~4~~ ~~5~~ ~~6~~ ~~7~~ ~~8~~ ~~9~~ ~~10~~ ~~11~~ ~~12~~ ~~13~~ ~~14~~ ~~15~~ ~~16~~ ~~17~~ ~~18~~ ~~19~~ ~~20~~ ~~21~~ ~~22~~ ~~23~~ ~~24~~ ~~25~~ ~~26~~ ~~27~~ ~~28~~ ~~29~~ ~~30~~ ~~31~~ ~~32~~ ~~33~~ ~~34~~ ~~35~~ ~~36~~ ~~37~~ ~~38~~ ~~39~~ ~~40~~ ~~41~~ ~~42~~ ~~43~~ ~~44~~ ~~45~~ ~~46~~ ~~47~~ ~~48~~ ~~49~~ ~~50~~ ~~51~~ ~~52~~ ~~53~~ ~~54~~ ~~55~~ ~~56~~ ~~57~~ ~~58~~ ~~59~~ ~~60~~ ~~61~~ ~~62~~ ~~63~~ ~~64~~ ~~65~~ ~~66~~ ~~67~~ ~~68~~ ~~69~~ ~~70~~ ~~71~~ ~~72~~ ~~73~~ ~~74~~ ~~75~~ ~~76~~ ~~77~~ ~~78~~ ~~79~~ ~~80~~ ~~81~~ ~~82~~ ~~83~~ ~~84~~ ~~85~~ ~~86~~ ~~87~~ ~~88~~ ~~89~~ ~~90~~ ~~91~~ ~~92~~ ~~93~~ ~~94~~ ~~95~~ ~~96~~ ~~97~~ ~~98~~ ~~99~~ ~~100~~ ~~101~~ ~~102~~ ~~103~~ ~~104~~ ~~105~~ ~~106~~ ~~107~~ ~~108~~ ~~109~~ ~~110~~ ~~111~~ ~~112~~ ~~113~~ ~~114~~ ~~115~~ ~~116~~ ~~117~~ ~~118~~ ~~119~~ ~~120~~ ~~121~~ ~~122~~ ~~123~~ ~~124~~ ~~125~~ ~~126~~ ~~127~~ ~~128~~ ~~129~~ ~~130~~ ~~131~~ ~~132~~ ~~133~~ ~~134~~ ~~135~~ ~~136~~ ~~137~~ ~~138~~ ~~139~~ ~~140~~ ~~141~~ ~~142~~ ~~143~~ ~~144~~ ~~145~~ ~~146~~ ~~147~~ ~~148~~ ~~149~~ ~~150~~ ~~151~~ ~~152~~ ~~153~~ ~~154~~ ~~155~~ ~~156~~ ~~157~~ ~~158~~ ~~159~~ ~~160~~ ~~161~~ ~~162~~ ~~163~~ ~~164~~ ~~165~~ ~~166~~ ~~167~~ ~~168~~ ~~169~~ ~~170~~ ~~171~~ ~~172~~ ~~173~~ ~~174~~ ~~175~~ ~~176~~ ~~177~~ ~~178~~ ~~179~~ ~~180~~ ~~181~~ ~~182~~ ~~183~~ ~~184~~ ~~185~~ ~~186~~ ~~187~~ ~~188~~ ~~189~~ ~~190~~ ~~191~~ ~~192~~ ~~193~~ ~~194~~ ~~195~~ ~~196~~ ~~197~~ ~~198~~ ~~199~~ ~~200~~ ~~201~~ ~~202~~ ~~203~~ ~~204~~ ~~205~~ ~~206~~ ~~207~~ ~~208~~ ~~209~~ ~~210~~ ~~211~~ ~~212~~ ~~213~~ ~~214~~ ~~215~~ ~~216~~ ~~217~~ ~~218~~ ~~219~~ ~~220~~ ~~221~~ ~~222~~ ~~223~~ ~~224~~ ~~225~~ ~~226~~ ~~227~~ ~~228~~ ~~229~~ ~~230~~ ~~231~~ ~~232~~ ~~233~~ ~~234~~ ~~235~~ ~~236~~ ~~237~~ ~~238~~ ~~239~~ ~~240~~ ~~241~~ ~~242~~ ~~243~~ ~~244~~ ~~245~~ ~~246~~ ~~247~~ ~~248~~ ~~249~~ ~~250~~ ~~251~~ ~~252~~ ~~253~~ ~~254~~ ~~255~~ ~~256~~ ~~257~~ ~~258~~ ~~259~~ ~~260~~ ~~261~~ ~~262~~ ~~263~~ ~~264~~ ~~265~~ ~~266~~ ~~267~~ ~~268~~ ~~269~~ ~~270~~ ~~271~~ ~~272~~ ~~273~~ ~~274~~ ~~275~~ ~~276~~ ~~277~~ ~~278~~ ~~279~~ ~~280~~ ~~281~~ ~~282~~ ~~283~~ ~~284~~ ~~285~~ ~~286~~ ~~287~~ ~~288~~ ~~289~~ ~~290~~ ~~291~~ ~~292~~ ~~293~~ ~~294~~ ~~295~~ ~~296~~ ~~297~~ ~~298~~ ~~299~~ ~~300~~ ~~301~~ ~~302~~ ~~303~~ ~~304~~ ~~305~~ ~~306~~ ~~307~~ ~~308~~ ~~309~~ ~~310~~ ~~311~~ ~~312~~ ~~313~~ ~~314~~ ~~315~~ ~~316~~ ~~317~~ ~~318~~ ~~319~~ ~~320~~ ~~321~~ ~~322~~ ~~323~~ ~~324~~ ~~325~~ ~~326~~ ~~327~~ ~~328~~ ~~329~~ ~~330~~ ~~331~~ ~~332~~ ~~333~~ ~~334~~ ~~335~~ ~~336~~ ~~337~~ ~~338~~ ~~339~~ ~~340~~ ~~341~~ ~~342~~ ~~343~~ ~~344~~ ~~345~~ ~~346~~ ~~347~~ ~~348~~ ~~349~~ ~~350~~ ~~351~~ ~~352~~ ~~353~~ ~~354~~ ~~355~~ ~~356~~ ~~357~~ ~~358~~ ~~359~~ ~~360~~ ~~361~~ ~~362~~ ~~363~~ ~~364~~ ~~365~~ ~~366~~ ~~367~~ ~~368~~ ~~369~~ ~~370~~ ~~371~~ ~~372~~ ~~373~~ ~~374~~ ~~375~~ ~~376~~ ~~377~~ ~~378~~ ~~379~~ ~~380~~ ~~381~~ ~~382~~ ~~383~~ ~~384~~ ~~385~~ ~~386~~ ~~387~~ ~~388~~ ~~389~~ ~~390~~ ~~391~~ ~~392~~ ~~393~~ ~~394~~ ~~395~~ ~~396~~ ~~397~~ ~~398~~ ~~399~~ ~~400~~ ~~401~~ ~~402~~ ~~403~~ ~~404~~ ~~405~~ ~~406~~ ~~407~~ ~~408~~ ~~409~~ ~~410~~ ~~411~~ ~~412~~ ~~413~~ ~~414~~ ~~415~~ ~~416~~ ~~417~~ ~~418~~ ~~419~~ ~~420~~ ~~421~~ ~~422~~ ~~423~~ ~~424~~ ~~425~~ ~~426~~ ~~427~~ ~~428~~ ~~429~~ ~~430~~ ~~431~~ ~~432~~ ~~433~~ ~~434~~ ~~435~~ ~~436~~ ~~437~~ ~~438~~ ~~439~~ ~~440~~ ~~441~~ ~~442~~ ~~443~~ ~~444~~ ~~445~~ ~~446~~ ~~447~~ ~~448~~ ~~449~~ ~~450~~ ~~451~~ ~~452~~ ~~453~~ ~~454~~ ~~455~~ ~~456~~ ~~457~~ ~~458~~ ~~459~~ ~~460~~ ~~461~~ ~~462~~ ~~463~~ ~~464~~ ~~465~~ ~~466~~ ~~467~~ ~~468~~ ~~469~~ ~~470~~ ~~471~~ ~~472~~ ~~473~~ ~~474~~ ~~475~~ ~~476~~ ~~477~~ ~~478~~ ~~479~~ ~~480~~ ~~481~~ ~~482~~ ~~483~~ ~~484~~ ~~485~~ ~~486~~ ~~487~~ ~~488~~ ~~489~~ ~~490~~ ~~491~~ ~~492~~ ~~493~~ ~~494~~ ~~495~~ ~~496~~ ~~497~~ ~~498~~ ~~499~~ ~~500~~ ~~501~~ ~~502~~ ~~503~~ ~~504~~ ~~505~~ ~~506~~ ~~507~~ ~~508~~ ~~509~~ ~~510~~ ~~511~~ ~~512~~ ~~513~~ ~~514~~ ~~515~~ ~~516~~ ~~517~~ ~~518~~ ~~519~~ ~~520~~ ~~521~~ ~~522~~ ~~523~~ ~~524~~ ~~525~~ ~~526~~ ~~527~~ ~~528~~ ~~529~~ ~~530~~ ~~531~~ ~~532~~ ~~533~~ ~~534~~ ~~535~~ ~~536~~ ~~537~~ ~~538~~ ~~539~~ ~~540~~ ~~541~~ ~~542~~ ~~543~~ ~~544~~ ~~545~~ ~~546~~ ~~547~~ ~~548~~ ~~549~~ ~~550~~ ~~551~~ ~~552~~ ~~553~~ ~~554~~ ~~555~~ ~~556~~ ~~557~~ ~~558~~ ~~559~~ ~~560~~ ~~561~~ ~~562~~ ~~563~~ ~~564~~ ~~565~~ ~~566~~ ~~567~~ ~~568~~ ~~569~~ ~~570~~ ~~571~~ ~~572~~ ~~573~~ ~~574~~ ~~575~~ ~~576~~ ~~577~~ ~~578~~ ~~579~~ ~~580~~ ~~581~~ ~~582~~ ~~583~~ ~~584~~ ~~585~~ ~~586~~ ~~587~~ ~~588~~ ~~589~~ ~~590~~ ~~591~~ ~~592~~ ~~593~~ ~~594~~ ~~595~~ ~~596~~ ~~597~~ ~~598~~ ~~599~~ ~~600~~ ~~601~~ ~~602~~ ~~603~~ ~~604~~ ~~605~~ ~~606~~ ~~607~~ ~~608~~ ~~609~~ ~~610~~ ~~611~~ ~~612~~ ~~613~~ ~~614~~ ~~615~~ ~~616~~ ~~617~~ ~~618~~ ~~619~~ ~~620~~ ~~621~~ ~~622~~ ~~623~~ ~~624~~ ~~625~~ ~~626~~ ~~627~~ ~~628~~ ~~629~~ ~~630~~ ~~631~~ ~~632~~ ~~633~~ ~~634~~ ~~635~~ ~~636~~ ~~637~~ ~~638~~ ~~639~~ ~~640~~ ~~641~~ ~~642~~ ~~643~~ ~~644~~ ~~645~~ ~~646~~ ~~647~~ ~~648~~ ~~649~~ ~~650~~ ~~651~~ ~~652~~ ~~653~~ ~~654~~ ~~655~~ ~~656~~ ~~657~~ ~~658~~ ~~659~~ ~~660~~ ~~661~~ ~~662~~ ~~663~~ ~~664~~ ~~665~~ ~~666~~ ~~667~~ ~~668~~ ~~669~~ ~~670~~ ~~671~~ ~~672~~ ~~673~~ ~~674~~ ~~675~~ ~~676~~ ~~677~~ ~~678~~ ~~679~~ ~~680~~ ~~681~~ ~~682~~ ~~683~~ ~~684~~ ~~685~~ ~~686~~ ~~687~~ ~~688~~ ~~689~~ ~~690~~ ~~691~~ ~~692~~ ~~693~~ ~~694~~ ~~695~~ ~~696~~ ~~697~~ ~~698~~ ~~699~~ ~~700~~ ~~701~~ ~~702~~ ~~703~~ ~~704~~ ~~705~~ ~~706~~ ~~707~~ ~~708~~ ~~709~~ ~~710~~ ~~711~~ ~~712~~ ~~713~~ ~~714~~ ~~715~~ ~~716~~ ~~717~~ ~~718~~ ~~719~~ ~~720~~ ~~721~~ ~~722~~ ~~723~~ ~~724~~ ~~725~~ ~~726~~ ~~727~~ ~~728~~ ~~729~~ ~~730~~ ~~731~~ ~~732~~ ~~733~~ ~~734~~ ~~735~~ ~~736~~ ~~737~~ ~~738~~ ~~739~~ ~~740~~ ~~741~~ ~~742~~ ~~743~~ ~~744~~ ~~745~~ ~~746~~ ~~747~~ ~~748~~ ~~749~~ ~~750~~ ~~751~~ ~~752~~ ~~753~~ ~~754~~ ~~755~~ ~~756~~ ~~757~~ ~~758~~ ~~759~~ ~~760~~ ~~761~~ ~~762~~ ~~763~~ ~~764~~ ~~765~~ ~~766~~ ~~767~~ ~~768~~ ~~769~~ ~~770~~ ~~771~~ ~~772~~ ~~773~~ ~~774~~ ~~775~~ ~~776~~ ~~777~~ ~~778~~ ~~779~~ ~~780~~ ~~781~~ ~~782~~ ~~783~~ ~~784~~ ~~785~~ ~~786~~ ~~787~~ ~~788~~ ~~789~~ ~~790~~ ~~791~~ ~~792~~ ~~793~~ ~~794~~ ~~795~~ ~~796~~ ~~797~~ ~~798~~ ~~799~~ ~~800~~ ~~801~~ ~~802~~ ~~803~~ ~~804~~ ~~805~~ ~~806~~ ~~807~~ ~~808~~ ~~809~~ ~~810~~ ~~811~~ ~~812~~ ~~813~~ ~~814~~ ~~815~~ ~~816~~ ~~817~~ ~~818~~ ~~819~~ ~~820~~ ~~821~~ ~~822~~ ~~823~~ ~~824~~ ~~825~~ ~~826~~ ~~827~~ ~~828~~ ~~829~~ ~~830~~ ~~831~~ ~~832~~ ~~833~~ ~~834~~ ~~835~~ ~~836~~ ~~837~~ ~~838~~ ~~839~~ ~~840~~ ~~841~~ ~~842~~ ~~843~~ ~~844~~ ~~845~~ ~~846~~ ~~847~~ ~~848~~ ~~849~~ ~~850~~ ~~851~~ ~~852~~ ~~853~~ ~~854~~ ~~855~~ ~~856~~ ~~857~~ ~~858~~ ~~859~~ ~~860~~ ~~861~~ ~~862~~ ~~863~~ ~~864~~ ~~865~~ ~~866~~ ~~867~~ ~~868~~ ~~869~~ ~~870~~ ~~871~~ ~~872~~ ~~873~~ ~~874~~ ~~875~~ ~~876~~ ~~877~~ ~~878~~ ~~879~~ ~~880~~ ~~881~~ ~~882~~ ~~883~~ ~~884~~ ~~885~~ ~~886~~ ~~887~~ ~~888~~ ~~889~~ ~~890~~ ~~891~~ ~~892~~ ~~893~~ ~~894~~ ~~895~~ ~~896~~ ~~897~~ ~~898~~ ~~899~~ ~~900~~ ~~901~~ ~~902~~ ~~903~~ ~~904~~ ~~905~~ ~~906~~ ~~907~~ ~~908~~ ~~909~~ ~~910~~ ~~911~~ ~~912~~ ~~913~~ ~~914~~ ~~915~~ ~~916~~ ~~917~~ ~~918~~ ~~919~~ ~~920~~ ~~921~~ ~~922~~ ~~923~~ ~~924~~ ~~925~~ ~~926~~ ~~927~~ ~~928~~ ~~929~~ ~~930~~ ~~931~~ ~~932~~ ~~933~~ ~~934~~ ~~935~~ ~~936~~ ~~937~~ ~~938~~ ~~939~~ ~~940~~ ~~941~~ ~~942~~ ~~943~~ ~~944~~ ~~945~~ ~~946~~ ~~947~~ ~~948~~ ~~949~~ ~~950~~ ~~951~~ ~~952~~ ~~953~~ ~~954~~ ~~955~~ ~~956~~ ~~957~~ ~~958~~ ~~959~~ ~~960~~ ~~961~~ ~~962~~ ~~963~~ ~~964~~ ~~965~~ ~~966~~ ~~967~~ ~~968~~ ~~969~~ ~~970~~ ~~971~~ ~~972~~ ~~973~~ ~~974~~ ~~975~~ ~~976~~ ~~977~~ ~~978~~ ~~979~~ ~~980~~ ~~981~~ ~~982~~ ~~983~~ ~~984~~ ~~985~~ ~~986~~ ~~987~~ ~~988~~ ~~989~~ ~~990~~ ~~991~~ ~~992~~ ~~993~~ ~~994~~ ~~995~~ ~~996~~ ~~997~~ ~~998~~ ~~999~~ ~~1000~~ ~~1001~~ ~~1002~~ ~~1003~~ ~~1004~~ ~~1005~~ ~~1006~~ ~~1007~~ ~~1008~~ ~~1009~~ ~~1010~~ ~~1011~~ ~~1012~~ ~~1013~~ ~~1014~~ ~~1015~~ ~~1016~~ ~~1017~~ ~~1018~~ ~~1019~~ ~~1020~~ ~~1021~~ ~~1022~~ ~~1023~~ ~~1024~~ ~~1025~~ ~~1026~~ ~~1027~~ ~~1028~~ ~~1029~~ ~~1030~~ ~~1031~~ ~~1032~~ ~~1033~~ ~~1034~~ ~~1035~~ ~~1036~~ ~~1037~~ ~~1038~~ ~~1039~~ ~~1040~~ ~~1041~~ ~~1042~~ ~~1043~~ ~~1044~~ ~~1045~~ ~~1046~~ ~~1047~~ ~~1048~~ ~~1049~~ ~~1050~~ ~~1051~~ ~~1052~~ ~~1053~~ ~~1054~~ ~~1055~~ ~~1056~~ ~~1057~~ ~~1058~~ ~~1059~~ ~~1060~~ ~~1061~~ ~~1062~~ ~~1063~~ ~~1064~~ ~~1065~~ ~~1066~~ ~~1067~~ ~~1068~~ ~~1069~~ ~~1070~~ ~~1071~~ ~~1072~~ ~~1073~~ ~~1074~~ ~~1075~~ ~~1076~~ ~~1077~~ ~~1078~~ ~~1079~~ ~~1080~~ ~~1081~~ ~~1082~~ ~~1083~~ ~~1084~~ ~~1085~~ ~~1086~~ ~~1087~~ ~~1088~~ ~~1089~~ ~~1090~~ ~~1091~~ ~~1092~~ ~~1093~~ ~~1094~~ ~~1095~~ ~~1096~~ ~~1097~~ ~~1098~~ ~~1099~~ ~~1100~~ ~~1101~~ ~~1102~~ ~~1103~~ ~~1104~~ ~~1105~~ ~~1106~~ ~~1107~~ ~~1108~~ ~~1109~~ ~~1110~~ ~~1111~~ ~~1112~~ ~~1113~~ ~~1114~~ ~~1115~~ ~~1116~~ ~~1117~~ ~~1118~~ ~~1119~~ ~~1120~~ ~~1121~~ ~~1122~~ ~~1123~~ ~~1124~~ ~~1125~~ ~~1126~~ ~~1127~~ ~~1128~~ ~~1129~~ ~~1130~~ ~~1131~~ ~~1132~~ ~~1133~~ ~~1134~~ ~~1135~~ ~~1136~~ ~~1137~~ ~~1138~~ ~~1139~~ ~~1140~~ ~~1141~~ ~~1142~~ ~~1143~~ ~~1144~~ ~~1145~~ ~~1146~~ ~~1147~~ ~~1148~~ ~~1149~~ ~~1150~~ ~~1151~~ ~~1152~~ ~~1153~~ ~~1154~~ ~~1155~~ ~~1156~~ ~~1157~~ ~~1158~~ ~~1159~~ ~~1160~~ ~~1161~~ ~~1162~~ ~~1163~~ ~~1164~~ ~~1165~~ ~~1166~~ ~~1167~~ ~~1168~~ ~~1169~~ ~~1170~~ ~~1171~~ ~~1172~~ ~~1173~~ ~~1174~~ ~~1175~~ ~~1176~~ ~~1177~~ ~~1178~~ ~~1179~~ ~~1180~~ ~~1181~~ ~~1182~~ ~~1183~~ ~~1184~~ ~~1185~~ ~~1186~~ ~~1187~~ ~~1188~~ ~~1189~~ ~~1190~~ ~~1191~~ ~~1192~~ ~~1193~~ ~~1194~~ ~~1195~~ ~~1196~~ ~~1197~~ ~~1198~~ ~~1199~~ ~~1200~~ ~~1201~~ ~~1202~~ ~~1203~~ ~~1204~~ ~~1205~~ ~~1206~~ ~~1207~~ ~~1208~~ ~~1209~~ ~~1210~~ ~~1211~~ ~~1212~~ ~~1213~~ ~~1214~~ ~~1215~~ ~~1216~~ ~~1217~~ ~~1218~~ ~~1219~~ ~~1220~~ ~~1221~~ ~~1222~~ ~~1223~~ ~~1224~~ ~~1225~~ ~~1226~~ ~~1227~~ ~~1228~~ ~~1229~~ ~~1230~~ ~~1231~~ ~~1232~~ ~~1233~~ ~~1234~~ ~~1235~~ ~~1236~~ ~~1237~~ ~~1238~~ ~~1239~~ ~~1240~~ ~~1241~~ ~~1242~~ ~~1243~~ ~~1244~~ ~~1245~~ ~~1246~~ ~~1247~~ ~~1248~~ ~~1249~~ ~~1250~~ ~~1251~~ ~~1252~~ ~~1253~~ ~~1254~~ ~~1255~~ ~~1256~~ ~~1257~~ ~~1258~~ ~~1259~~ ~~1260~~ ~~1261~~ ~~1262~~ ~~1263~~ ~~1264~~ ~~1265~~ ~~1266~~ ~~1267~~ ~~1268~~ ~~1269~~ ~~1270~~ ~~1271~~ ~~1272~~ ~~1273~~ ~~1274~~ ~~1275~~ ~~1276~~ ~~1277~~ ~~1278~~ ~~1279~~ ~~1280~~ ~~1281~~ ~~1282~~ ~~1283~~ ~~1284~~ ~~1285~~ ~~1286~~ ~~1287~~ ~~1288~~ ~~1289~~ ~~1290~~ ~~1291~~ ~~1292~~ ~~1293~~ ~~1294~~ ~~1295~~ ~~1296~~ ~~1297~~ ~~1298~~ ~~1299~~ ~~1300~~ ~~1301~~ ~~1302~~ ~~1303~~ ~~1304~~ ~~1305~~ ~~1306~~ ~~1307~~ ~~1308~~ ~~1309~~ ~~1310~~ ~~1311~~ ~~1312~~ ~~1313~~ ~~1314~~ ~~1315~~ ~~1316~~ ~~1317~~ ~~1318~~ ~~1319~~ ~~1320~~ ~~1321~~ ~~1322~~ ~~1323~~ ~~1324~~ ~~1325~~ ~~1326~~ ~~1327~~ ~~1328~~ ~~1329~~ ~~1330~~ ~~1331~~ ~~1332~~ ~~1333~~ ~~1334~~ ~~1335~~ ~~1336~~ ~~1337~~

media. Data confirmed that this skin reactivity was due to cholera toxin (see Finkelstein and LoSpallutto, 1969).

Craig (1965) cautioned, "The absence of skin lesions in clinical cholera certainly does not preclude the possibility that the noxa responsible for gut damage could also have a deleterious effect upon the skin provided it is applied to skin in sufficient concentration". The extreme reactogenicity of cholera toxin in the skin was used as a test for its toxicity and the prior art evidenced an expectation that cholera toxin would be reactogenic if applied to the skin, producing an undesirable reaction. Such adverse reactions have been well documented by known authorities in the field (Craig, 1972).

In contrast, we have shown cholera toxin to be immunogenic, acting as both antigen and adjuvant, when placed on the skin but without any resulting local or systemic side effects. This lack of reactogenicity when cholera toxin was placed on the skin for transcutaneous immunization was surprising and contradicted conclusions one would have drawn from the prior art. Specifically, CT placed on the skin according to our invention acts as a non-toxic, non-reactogenic adjuvant, in contrast to the expectations of Craig, while injection of CT into the skin results in swelling and redness. Thus, it was not obvious prior to our invention that cholera toxin or other ADP-ribosylating exotoxins or other adjuvants applied topically would be useful for transcutaneous immunization. In fact large doses of heat labile enterotoxin (LT) placed on the skin of humans has been shown to induce a systemic immune response without local or systemic toxicity.

This unexpected absence of reactogenicity is extremely important to the use of vaccines. Vaccine antigens and adjuvants are useful when immunization produces a protective immune response without significant unwanted reactions. Historically, reactogenicity of vaccines such as swelling, tenderness and pain at the site of injection has in some cases (e.g., typhoid and pertussis) been accepted because of the benefits of vaccination. However, high levels of reactogenicity and other side effects are not desirable, and would be problematic for development of new vaccine adjuvant and antigen candidates. Research efforts are focused on making vaccine adjuvants that are stimulatory and do not inducing unwanted reactions. Whole cell pertussis vaccines induce systemic and local side effects and, as a result, this effective vaccine and time

tested vaccine is being replaced by acellular pertussis vaccines solely because they are less reactogenic.

The present invention differs from that of Patent No. 5,830,877 which teaches the use of a naked plasmid that encodes for biologically active peptides into a mammalian host. The invention described herein teaches the use of an adjuvant and antigen or nucleic acid administered together on the skin to induce an immune response. The invention described herein further differs from that of Patent # 5,830,877 which teaches away from the use of peptides that are not encoded in a nucleic acid and produced by the host cell because of the toxicity associated with biologically active peptides, the problems and cost of isolating, purifying and synthesizing peptides and their short half life in vivo resulting from degradation by proteases present in the target tissue. This clearly teaches away from the addition of an adjuvant such as a cholera toxin to a coadministered antigen or nucleic acid. In fact the novelty of the ability of a large molecule such as CT to induce an immune response by application through the skin without toxicity has led to a number of scientific papers describing this novelty and public excitement over the potential implication of delivery of proteins for vaccination by skin application. Unlike Patent No. 5,830,877 the present invention is not dependent on stimulation of local inflammation or irritation. Unlike Patent No. 5,830,877 the invention does not depend on irritation or local inflammation to increase the permeability of cell membranes to enhance the uptake of the antigens, plasmids or RNA. In fact the striking feature regarding Transcutaneous Immunization is the absence of local inflammation.

Unlike the present invention, Patent No. 5,824,313 teaches the application of extremely small (less than 500 daltons) lymphoid organ modifying agents such as 1,25-dihydroxy-16-ene Vitamin D<sub>3</sub> and calcipotriene or dehydroepiandrosterone (DHEA), DHEA congeners and DHEA-derivatives with the intramuscular injection of an antigen to affect antibody responses.

Transcutaneous immunization requires both passage of an antigen through the outer barriers of the skin, which was thought to be impervious to such passage, and an immune response to the antigen. Fisher's Contact Dermatitis states that molecules of greater than 500 daltons cannot normally penetrate the skin. There is a report by Paul et al. (1995) of induction of an immune response with transferosomes, a lipid structure

distinct from liposomes. In this publication, the transferosomes were used as a vehicle for antigen (bovine serum albumin and gap junction proteins) and complement-mediated lysis of antigen-sensitized liposomes was assayed. The limit to penetration of the skin by antigen was stated to be 750 daltons. In their study, an immune response was not induced when a solution containing antigen was placed on the skin; only transferosomes were able to induce an immune response. Paul and Cvec (1995) also stated that it is "impossible to immunize epicutaneously with simple peptide or protein solutions".

Such references explain why our successful use of a molecule like cholera toxin (which is 85,000 daltons) as an antigen or adjuvant in immunization was greeted with surprise by the field because such large molecules were not expected to penetrate the skin and, therefore, would not be expected to induce a specific immune response.

However, we have shown in U.S. Appln. No. 08/749,164 (filed November 14, 1996); U.S. Appln. No. 08/896,085 (filed July 17, 1997); and international application PCT/US97/21324 (filed November 14, 1997) that using an ADP-ribosylating exotoxin, such as cholera toxin, as an antigen could elicit a strong antibody response that is highly reproducible. When an ADP-ribosylating exotoxin, such as cholera toxin, was used as an immunoadjuvant and applied to the skin in a saline solution with a separate antigen (e.g., diphtheria toxoid), a systemic and mucosal antigen-specific antibody response could be elicited. In the present application, we disclose that transcutaneous immunization using a penetration enhancer, a barrier disruption agent, or combinations thereof may improve the adjuvant activity of a bacterial exotoxin.

We have shown that like cholera toxin (CT), heat-labile enterotoxin from E. coli (LT), Pseudomonas exotoxin A (ETA), pertussis toxin (PT) and a wide variety of antigens including killed rabies virus, recombinants such as HIV p55 gag, polysaccharide conjugates such as Hib, sonicates, pertactin for example, are able to pass through the skin and induce an immune response. Additionally CT, LT, ETA and PT and bacterial DNA and cytobines, can act as adjuvants to induce an immune response to antigens co-administered on the skin. Thus tetanus toxoid, not immunogenic by itself on the skin, can induce a strong immune response when placed on the skin with CT. We have proposed that the Langerhans cell population underlying the site of application are a preferred antigen presenting cell for delivering

antigen to the immune system. Adjuvant may act on the antigen presenting cell directly, or through lymphocytes recognizing antigen.

We propose to enhance the immune response to transcutaneously adjuvant and/or antigen utilizing penetration enhancement techniques. According to Hurley,  
5 "Skin owes its durability to the dermis, but its chemical impermeability resides in the epidermis and almost exclusively in its dead outer layer, the stratum corneum". For transcutaneous immunization using, for example, an ADP-ribosylating exotoxin as adjuvant and a soluble protein antigen such as diphtheria toxoid, penetration of the stratum corneum must occur. Penetration enhancement techniques would be designed  
10 to increase the movement of transcutaneous antigens and adjuvants through the stratum corneum layer of skin.

Further, we propose that transcutaneous immunization using activation of at least one of the antigen, adjuvant and skin component will enhance the immune response as assayed by quantitative and qualitative parameters. Antigen-adjuvant of  
15 the formulation may be activated by trypsin cleavage of a bacterial exotoxin (e.g., trypsin-cleaved LT, with or without reduction). Activation of the skin at the application site of the formulation may be accomplished by the use of barrier disruption agents (e.g., acetone, alcohol) which increases the size or activation of the underlying Langerhans cell population, or by an enzyme or combination of enzymes  
20 (e.g., enzymes with sialidase activity) which increases the amount or accessibility of ganglioside GM1 receptor.

## SUMMARY OF THE INVENTION

25 An object of the invention is to provide an enhanced system for transcutaneous immunization which induces an immune response (e.g., humoral and/or cellular effector) in a subject, a subject being an animal or human. This delivery system provides simple application to intact skin of an organism of a formulation comprised of antigen and adjuvant to induce a specific immune response  
30 against the antigen. Although not required for induction of an immune response by this simple delivery system, supplementation of the aforementioned process with penetration enhancement or barrier disruption may enhance immunization and/or vaccination.

In particular, the adjuvant or antigen or skin may assist in the penetration of the stratum corneum or epidermis to encounter the antigen presenting cells of the immune system (e.g., Langerhans cells in the epidermis, dermal dendritic cells, dendritic cells, follicular dendritic cells, macrophages, B lymphocytes) and/or induce the antigen presenting cells to phagocytose the antigen. The antigen presenting cells then present the antigen to T and B cells. In the instance of Langerhans cells, the antigen presenting cells then may migrate from the skin to the lymph nodes and present antigen to lymphocytes (e.g., B and/or T cells), thereby inducing an antigen-specific immune response.

In addition activation of the antigen, adjuvant, skin, or any combination thereof may be accomplished to supplement the immunization process.

In addition to eliciting immune reactions leading to generation of an antigen-specific B lymphocyte and/or T lymphocyte, including a cytotoxic T lymphocyte (CTL), another object of the invention is to positively and/or negatively regulate components of the immune system by using the transcutaneous immunization system to affect antigen-specific helper (Th1 and/or Th2) or delayed-type hypersensitivity (DTH) T-cell subsets. This can be exemplified by the differential behavior of CT and LT which can result in different T-helper responses or different levels of protection in in-vivo challenge molds using transcutaneous immunization.

## DESCRIPTION OF THE DRAWINGS

Figures 1a-f are photographs showing no inflammation at the site of immunization (A,B), Langerhans cell activation by LT in human skin at the site of immunization (C,E) and the absence of Langerhans cell activation in skin from the contralateral arm (D,F).

Figures 2a-d are photographs showing normal Langerhans cell (A,B, 200 and 400X) and Langerhan cell activation by Cholera Toxin in mouse skin (C,D. 200X, 400X).



## DESCRIPTION OF PREFERRED EMBODIMENTS

In one embodiment of the invention, a formulation containing antigen and adjuvant such as CT and DT, is applied to intact skin of an organism after penetration enhancement of the skin, the antigen is presented to immune cells, and an antigen-specific immune response is induced without perforating the skin. The formulation may include additional antigens or nucleic acids such that transcutaneous application of the formulation induces an immune response to multiple antigens, or nucleic acids encoding for antigens preferably from 2 to 20 but possibly up to 200. In such a case, the antigens may or may not be derived from the same source, but the antigens will have different chemical structures so as to induce immune responses specific for the different antigens. Antigen-specific lymphocytes may participate in the immune response and, in the case of participation by B lymphocytes, antigen-specific antibodies may be part of the immune response.

In another embodiment of the invention, the invention is used to treat an organism. If the antigen is derived from a pathogen, the treatment vaccinates the organism against infection by the pathogen or against its pathogenic effects such as those caused by toxin secretion. A formulation that includes a tumor antigen may provide a cancer treatment; a formulation that includes an allergen may be used to treat for allergic disease; a formulation that includes an autoantigen may provide a treatment for a disease caused by the organism's own immune system (e.g., autoimmune disease). The invention may be used therapeutically to treat existing disease, protectively to prevent disease, or to reduce the severity and/or duration of disease.

In a further embodiment of the invention, a patch for use in the above methods is provided. The patch may comprise a dressing, and effective amounts of antigen or nucleic acids and adjuvant. The dressing may be occlusive or non-occlusive. The patch may contain penetration enhancers or may include a device for physical penetration enhancement. The patch may include additional antigens such that application of the patch induces an immune response to multiple antigens. In such a case, the antigens may or may not be derived from the same source, but the antigens will have different chemical structures so as to induce an immune response specific

for the different antigens. For effective treatment, multiple patches may be applied at frequent intervals or constantly over a period of time.

Moreover, in yet another embodiment of the invention, the formulation is applied to intact skin overlying more than one draining lymph node field using either single or multiple applications or a separate patch for adjuvant or antigen/nucleic acid. The formulation may include additional antigens such that application to intact skin induces an immune response to multiple antigens. In such a case, the antigens may or may not be derived from the same source, but the antigens will have different chemical structures so as to induce an immune response specific for the different antigens.

The formulation may be applied to the skin to boost or prime the immune response in conjunction with other routes of immunization. Thus, priming with transcutaneous immunization with either single or multiple applications may be followed with oral, nasal, or parenteral techniques for boosting immunization with the same or altered antigens. The formulation may include additional antigens such that application to intact skin induces an immune response to multiple antigens. In such a case, the antigens may or may not be derived from the same source, but the antigens will have different chemical structures so as to induce an immune response specific for the different antigens.

In addition to antigen and activated adjuvant, the formulation may comprise a vehicle. For example, the formulation may comprise AQUAPHOR (an emulsion of petrolatum, mineral oil, mineral wax, wool wax, panthenol, bisabol, and glycerin), emulsions (e.g., aqueous creams), microemulsions, gels, oil-in-water emulsions (e.g., oily creams), anhydrous lipids and oil-in-water emulsions, anhydrous lipids and water-in-oil emulsions, fats, waxes, oil, silicones, and humectants (e.g., glycerol).

The antigen may be derived from a pathogen that can infect the organism (e.g., bacterium, virus, fungus, or parasite), or a cell (e.g., tumor cell or normal cell) or allergen or biological warfare agent. The antigen may be a tumor antigen or an autoantigen. Chemically, the antigen may be a carbohydrate, glycolipid, glycoprotein, lipid, lipoprotein, phospholipid, polypeptide, or fusion protein (recombinant) or chemical conjugate of the above. The molecular weight of the antigen may be greater than 500 daltons, preferably greater than 800 daltons, and more preferably greater than 1000 daltons.

Antigen may be obtained by recombinant means, chemical synthesis, or purification from a natural source. One advantage of transcutaneous immunization may be that purification of an antigen is not necessary e.g. a whole organism may be sonicated and used for immunization. The level of toxicity associated with injecting a product from such a preparation is often too toxic to be tolerated, such as LPS, which can be fatal if injected, but is non-toxic on the skin. Preferred are proteinaceous antigen or conjugates with polysaccharide. Antigen may be at least partially purified in cell-free form. Alternatively, antigen may be provided in the form of a live virus, an attenuated live virus, or an inactivated virus, sonicated or lysed whole bacterium, parasite or detergent treated virus, or fraction thereof.

Inclusion of an adjuvant may allow potentiation or modulation of the immune response. Moreover, selection of a suitable antigen or adjuvant may allow preferential induction of a humoral or cellular immune pr mucosal response, specific antibody isotypes (e.g., IgM, IgD, IgA1, IgA2, IgE, IgG1, IgG2, IgG3, and/or IgG4), and/or specific T-cell subsets (e.g., CTL, Th1, Th2 and/or T<sub>DTH</sub>). Optionally, antigen, adjuvant, may be provided in the formulation by means of a nucleic acid (e.g., DNA, RNA, cDNA, cRNA) encoding the antigen or adjuvant as appropriate with a antigen or adjuvant that has been added to the nucleic acid. This technique is called genetic immunization.

The term "antigen" as used in the invention, is meant to describe a substance that induces a specific immune response when presented to immune cells of an organism. An antigen may comprise a single immunogenic epitope, or a multiplicity of immunogenic epitopes recognized by a B-cell receptor (i.e., antibody on the membrane of the B cell) or a T-cell receptor. A molecule may be both an antigen and an adjuvant (e.g., cholera toxin) and, thus, the formulation may contain only one component. Antigen may be provided as a whole organism such as, for example, a bacterium or virion; antigen may be obtained from an extract or lysate, either from whole cells or membrane alone; or antigen may be chemically synthesized for produced by recombinant means.

The term "adjuvant" as used in the invention, is meant to describe a substance added to the formulation to assist in inducing an immune response to the antigen.

The term "effective amount" as used in the invention, is meant to describe that amount of antigen which induces an antigen-specific immune response. Such

induction of an immune response may provide a treatment such as, for example, immunoprotection, desensitization, immunosuppression, modulation of autoimmune disease, potentiation of cancer immunosurveillance, or therapeutic vaccination against an established infectious disease.

5 By the epidermis we mean the cells of the skin from the basal layer of kertinoctyes and basal lamina up to and through the stratum corneum.

The definition of transdermal is generally held to be: Relating to, being, or supplying a medication in a form for absorption through the skin into the bloodstream (~drug delivery) (~nitroglycerin) (~nicotine patch). <sup>2</sup>Frederick C. Mish et al., eds.,  
10 *Merriam-Webster's Collegiate Dictionary*, 10<sup>th</sup> ed. (Springfield, MA.: Merriam-Webster, Incorporated, 1997), 861.

The term draining lymph node field as used in the invention means an anatomic area over which the lymph collected is filtered through a set of defined set of lymph nodes (e.g., cervical, axillary, inguinal, epitrochelear, popliteal, those of the  
15 abdomen and thorax).

Skin penetration may be enhanced using techniques that increase skin hydration. According to Roberts and Walker (1993), "The state of hydration of the stratum corneum (SC) is one of the most important factors in determining the rate of percutaneous absorption of a given solute". The state of hydration interacts with the  
20 principal of diffusion in determining the rate of absorption of a substance through the skin. Furthermore, Hurley stated:

"Absorption of substances through the stratum corneum is believed to occur by diffusion in accordance with Fick's laws of diffusion in which the rate of absorption of a chemical is proportional to the  
25 concentration difference across the membrane. Thus, a concentration gradient between the high concentration of solute on the skin surface and its absence or low concentration below the stratum corneum is the driving force in this process. Transcorneal movement of absorption is classically represented as 'percellular', that is, directly through the cell  
30 walls of the compacted corneum and not intercellularly. Intracellular protein filaments are described as the pathways for polar (water soluble) compounds and the medium between filaments serves as the route for nonpolar (lipid-soluble) substances. . . Hydration increases the permeability of the stratum corneum for most substances by a  
35 number of cytophysical mechanisms that are not completely clarified."

Thus, while skin hydration is generally known to enhance skin penetration, the mechanisms by which this occurs are not entirely clear and, thus, not predictable prior

to the present invention and not thought to allow penetration of large molecules (7750 daltons).

The use of vehicles for increasing hydration is well known. Occlusive dressings, such as vapor-impenetrable plastic films (e.g., polyvinylidene, polyethylene) enhance absorption principally through increased hydration of the stratum corneum, a result of swelling of the corneocytes, and uptake of water into the intercellular corridors. Hydrocolloid patches may also be used to enhance skin penetration. Absorption of steroids can be increased over 100 fold using plastic occlusive film. Generally, greases, oils or impermeable plastic induce the most hydration by occlusion. See, for example, Idson (1978); Hollingsbee (1995); and McKenzie and Stoughton (1962). The use of hydration or vehicle for hydration with an antigen and adjuvant were not known prior to our invention as penetration. The skin was thought to be limited to small molecules even in the hydrated state.

Suitable agents which are known to enhance absorption of drugs through skin are described in Sloan, Use of Solubility Parameters from Regular Solution Theory to Describe Partitioning-Driven Processes, Ch. 5, "Prodrugs: Topical and Ocular Drug Delivery" (Marcel Dekker, 1992), and at places elsewhere in the text.

It is expected that these techniques (and others which are conventionally used to facilitate drug delivery) may be adapted to preparation of nucleic acids for use in the methods of the invention by those of ordinary skill in the art without undue experimentation. Specific examples illustrating this suitability are set forth below.

The state of the art in skin penetration enhancement is described in Pharmaceutical Skin Penetration Enhancement Edited by Kenneth A. Walters and Jonathan Hadgraft, Published by Marcel Dekker, Inc., New York, 1993.

Skin permeability and/or skin hydration may be expected by selecting an appropriate vehicle from different classes, such as humectant (e.g., glycols, glycerols), powder, (e.g., clays, shake lotions), oil/water (O/W) emulsion (e.g., aqueous creams), water/oil emulsion (e.g., oily creams), emulsifying base (e.g., anhydrous lipid and O/W emulsifiers), absorption base (e.g., anhydrous lipid and W/O emulsifiers), lipophilic (e.g., fats, waxes, oils, silicones), and occlusive dressing (e.g., plastic wrap).

Other methods that disrupt the stratum corneum proteins to enhance penetration in the present invention may be employed. Salicylic acid is a keratinolytic that may increase absorption. Urea acts both as a keratinolytic and hydrater of the

skin, and may act as a penetration enhancer. Phospholipase A2 and phosphatidylcholine dependent phospholipase C may be used as epidermal enzymes to enhance penetration. Other penetration enhancers may include ethanol, acetone, detergents, bases, nair®, propylene glycol, pyrrolidones, dimethylacetamide, dimethylformamide, dimethylsulfoxide, alkyl sulfoxide, phosphine oxide, surfactants and caprolactams such as azone. Other compounds that may be used for penetration enhancement include amines and amides, alkyl N,N-distributed-amino acetates, decylmethylsulfoxide, pyrrolidones, pirotidecane (HPE-101), benzylalkonium, benzylalkonium chloride polymers, silicone based polymers, fatty acids, cyclic ureas, terpenes, liposomes, and cyclodextrins. Penetration enhancers are well known in the art, for example as described in Pharmaceutical Penetration Enhancement, (Marcel Dekker, 1993) Other techniques that may be employed for penetration include iontophoresis, ultrasound, electroporation, tape stripping, the use of gene guns or other propellant devices, tines such as used for TB tine tests (as provided by Mono-Vacc system) or microneedles which penetrate the outer surface of the skin, or abrasives which remove the outer layers of the skin and lipid extraction.

A device which may be used for disruption of the stratum corneum (which is distributed in the U.S. by Connaught Laboratories, Inc. of Swiftwater, Pa.) consists of a plastic container having a syringe plunger at one end and a tyne disk at the other. The tyne disk supports a multiplicity of narrow diameter tynes of a length which will just scratch the outermost layer of epidermal cells but not penetrate the epidermis. Each of the tynes in the MONO-VACC kit is coated with old tuberculin; in the present invention, each needle may be coated with a pharmaceutical composition of antigen/nucleic acid and adjuvant. Use of the device with the present invention may not be according to the manufacturer's written instructions included with the device product because when used with the present invention does not penetrate the epidermis. Hereto, the device may be used for surface disruption to disrupt the outermost layers of the skin, the stratum corneum and upper epidermis, to enhance the transcutaneous immunization. Similar devices which may also be used in this embodiment are those which are currently used to perform allergy tests.

Other approaches include barrier disruption. Inhibition of cholesterol synthesis using systemically administered HMG CoA reductase inhibitors and similar

drugs interfere with barrier function and may allow enhanced penetration of the formulation component.

It is also conceivable that the skin can be transformed to enhance the transcutaneous immune response. CT and LT exert their effect via the ganglioside GM1 binding by the B subunit. Ganglioside GM1 is a ubiquitous cell membrane glycolipid found in all mammalian cells. In the gastrointestinal tract, when the pentameric CT B subunit binds to the cell surface, a hydrophilic pore is formed which allows the A subunit to penetrate across the lipid bilayer. The skin contains gangliosides at a concentration of 30-35 nmol NeuAC/gm. Skin gangliosides are possible targets for initiating transcutaneous immunization via mechanisms such as Langerhans cells activation as described above.

One possible method to activate the skin for enhancing the effect of transcutaneous immunization with ADP ribosylating exotoxins by increasing the number of GM1 ganglioside molecules in skin cells. This could be achieved by activation of receptor cells using sialidase to convert gangliosides that do not bind the toxin into the sialidase-stable cholera toxin binding ganglioside GGnSLC (ganglioside GM1):

“It is interesting that the cholera vibrio is perhaps the best-known source of sialidase (or neuraminidase, as it is often called). Could this sialidase play a part in the natural history of the disease by making more receptors available for the toxin? If so, should any active immunizing agent the disease contain an anti-neuriminidase element? Incubation of intestinal scrapings with sialidase leads to a considerable increase in their ability to bind the toxin, which is due not only to conversion of sialidase-labile ganglioside to cholera toxin-binding ganglioside, but also, apparently to the unmasking of otherwise unapproachable ganglioside binding sites possibly by breaking down glycoproteins. Pretreatment of dog intestine with sialidase makes it produce more fluid in response to cholera toxin; treatment of adrenal cells with sialidase increases their responsiveness to cholera toxin; pretreatment of pigeon red cells with sialidase increases the activation of the adenylate cyclase in them by cholera toxin.

The biochemistry of cholera, In: Cholera: The American Scientific Experience, 1947-1980, van Heyningen, W.E., and Seal, J.R., Eds, Waterview Press, Boulder, 1983, page 263 (citations omitted).

The effect of treatment of the skin with sialidase may enhance the binding of an ADP ribosylating exotoxin such as CT to the immune cells targeted by transcutaneous immunization. This represents a kind of activation of the skin for transcutaneous immunization. Additionally, neuraminidase may act as an epidermal enzyme concurrently enhancing penetration.

The use of a penetration enhancer may be used in conjunction with activation of the skin. Activation of the skin for transcutaneous immunization may also be after treatments such as acetone or alcohol swabbing. It has been shown that skin barrier disruption using acetone swabbing of the skin increased Langerhans cell density by 80% and increased the reaction to contact allergens in vivo. If the density of Langerhans cells is increased, then the potency of the immune response may be increased. Similar chemical disruption might be expected to increase the number of Langerhans cells and result in activation of the skin component of transcutaneous immunization, by tape stripping, sodium dodecyl sulfate, the use of alcohol swabbing, or a depilatory such as calcium hydroxide. See Proksch and Brasch (1996, 1997) for use of penetration enhancers and barrier disruption in allergic contact dermatitis.

Penetration enhancement may be achieved by performance of simple maneuvers such as alcohol swabbing immediately prior to immunization, by concurrent use of penetration enhancement compounds or techniques or by techniques such as acetone swabbing 24 hours prior to increase the number of Langerhans cells.

Processes for preparing a pharmaceutical formulation are well-known in the art, whereby the antigen and adjuvant is combined with a pharmaceutically acceptable carrier vehicle. Suitable vehicles and their preparation are described, for example, in Remington's Pharmaceutical Sciences by E.W. Martin. Such formulations will contain an effective amount of the antigen and adjuvant together with a suitable amount of vehicle in order to prepare pharmaceutically acceptable compositions suitable for administration to a human or animal. The formulation may be applied in the form of an cream, emulsion, gel, lotion, ointment, paste, solution, suspension, or other forms known in the art. In particular, formulations that enhance skin hydration, penetration, or both are preferred. There may also be incorporated other pharmaceutically acceptable additives including, for example, diluents, excipients, binders, stabilizers, preservatives, and colorings.



Without being bound to any particular theory but only to provide an explanation for our observations, it is presumed that the transcutaneous immunization delivery system carries antigen to cells of the immune system where an immune response is induced. The antigen may pass through the normal protective outer layers of the skin (i.e., stratum corneum) and induce the immune response directly, or through an antigen presenting cell (e.g., macrophage, tissue macrophage, Langerhans cell, dendritic cell, dermal dendritic cell, B lymphocyte, or Kupffer cell) that presents processed antigen to a T lymphocyte (see Stingl et al., 1989; Streilein and Grammer, 1989; Tew et al., 1997). Optionally, the antigen may pass through the stratum corneum via a hair follicle or a skin organelle (e.g., sweat gland, oil gland).

Transcutaneous immunization with bacterial ADP-ribosylating exotoxins (bAREs) may target the epidermal Langerhans cell, known to be among the most efficient of the antigen presenting cells (APCs). We have found that bAREs activate Langerhans cells when applied epicutaneously to the skin in saline solution.

Adjuvants such as activated LT may greatly enhance Langerhans cell activation. The Langerhans cells direct specific immune responses through phagocytosis of the antigens, and migration to the lymph nodes where they act as APCs to present the antigen to lymphocytes, and thereby induce a potent antibody response. Although the skin is generally considered a barrier to invading organisms, the imperfection of this barrier is attested to by the numerous Langerhans cells distributed throughout the epidermis that are designed to orchestrate the immune response against organisms invading via the skin. According to Udey (1997):

"Langerhans cells are bone-marrow derived cells that are present in all mammalian stratified squamous epithelia. They comprise all of the accessory cell activity that is present in uninflamed epidermis, and in the current paradigm are essential for the initiation and propagation of immune responses directed against epicutaneously applied antigens. Langerhans cells are members of a family of potent accessory cells ('dendritic cells') that are widely distributed, but infrequently represented, in epithelia and solid organs as well as in lymphoid tissue.

"It is now recognized that Langerhans cells (and presumably other dendritic cells) have a life cycle with at least two distinct stages. Langerhans cells that are located in epidermis constitute a regular network of antigen-trapping 'sentinel' cells. Epidermal Langerhans cells can ingest particulates, including microorganisms, and are efficient processors of complex antigens. However, they express only low levels of MHC class I and II antigens and costimulatory molecules

(ICAM-1, B7-1 and B7-2) and are poor stimulators of unprimed T cells. After contact with antigen, some Langerhans cells become activated, exit the epidermis and migrate to T-cell-dependent regions of regional lymph nodes where they local as mature dendritic cells. In the course of exiting the epidermis and migrating to lymph nodes, antigen-bearing epidermal Langerhans cells (now the 'messengers') exhibit dramatic changes in morphology, surface phenotype and function. In contrast to epidermal Langerhans cells, lymphoid dendritic cells are essentially non-phagocytic and process protein antigens inefficiently, but express high levels of MHC class I and class II antigens and various costimulatory molecules and are the most potent stimulators of naive T cells that have been identified."

We envision that the potent antigen presenting capability of the epidermal Langerhans cells can be exploited for transcutaneously delivered vaccines. A transcutaneous immune response using the skin immune system would require delivery of vaccine antigen only to Langerhans cells in the stratum corneum (the outermost layer of the skin consisting of cornified cells and lipids) via passive diffusion and subsequent activation of the Langerhans cells to take up antigen, migrate to B-cell follicles and/or T-cell dependent regions, and present the antigen to B and/or T cells. If antigens other than bAREs (for example diphtheria toxoid) are to be phagocytosed by the Langerhans cells, then these antigens could also be taken to the lymph node for presentation to T-cells and subsequently induce an immune response specific for that antigen (e.g., diphtheria toxoid). Thus, a feature of transcutaneous immunization is the activation of the Langerhans cell, presumably by bacterial ADP-ribosylating exotoxins, ADP-ribosylating exotoxin binding subunits (e.g., cholera toxin B subunit), or other adjuvants or Langerhans cell activating substance. Increasing the skin population of Langerhans cells using strategies such as acetone swabbing could then be expected to enhance the transcutaneous immune response.

The spectrum of more commonly known skin immune responses is represented by contact dermatitis and atopy. Contact dermatitis, a pathogenic manifestation of LC activation, is directed by Langerhans cells which phagocytose antigen, migrate to lymph nodes, present antigen, and sensitize T cells that migrate to the skin and cause the intense destructive cellular response that occurs at affected skin sites (Dahl, 1996; Leung, 1997). Atopic dermatitis may utilize the Langerhans cell in a similar fashion, but is identified with Th2 cells and is generally associated with high levels of IgE antibody (Dahl, 1996; Leung, 1997).

Transcutaneous immunization with cholera toxin and related bAREs on the other hand is a novel immune response with an absence of superficial and microscopic post-immunization skin findings (i.e., non-inflamed skin) shown by the absence of lymphocyte infiltration 24, 48 and 120 hours after immunization. This is strikingly shown by completion of a Phase I trial in which humans were immunized with LT under a simple occlusive patch. Potent anti-LT IgG and IgA antibodies were stimulated. Two volunteers had biopsies performed at the site of immunization. Micro-scope evaluation confirmed the clinical observation that no inflammation was seen. This suggests that Langerhans cells, which "comprise all of the accessory cell activity that is present in uninflamed epidermis, and in the current paradigm are essential for the initiation and propagation of immune responses directed against epicutaneously applied antigens" (Udey, 1997) may have been recruited. The uniqueness of the transcutaneous immune response here is also indicated by the both high levels of antigen-specific IgG antibody, and the type of antibody produced (e.g., IgG1, IgG2a, IgG2b, IgG3 and IgA) and the absence of anti-CT IgE antibody. However, other immune cells may be engaged and speculation on the mechanism should not limit the invention.

Thus, we have found that bacterial-derived toxins applied to the surface of the skin can activate Langerhans cells and that TCI induces a potent immune response manifested as high levels of antigen-specific circulating IgG antibodies and would expect that penetration enhancement would enhance the immune response. Transcutaneous adjuvant and penetration enhancer may be used in transcutaneous immunization to enhance the IgG antibody or T-cell response to proteins not otherwise immunogenic by themselves when placed on the skin.

Transcutaneous targeting of Langerhans cells may also be used to deactivate their antigen presenting function, thereby preventing immunization or sensitization. Techniques to mobilize Langerhans cells or other skin immune cells yet negatively modulate them include, for example, the use of anti-inflammatory steroidal or non-steroidal agents (NSAID), cyclophosphamide or other immunosuppressants, interleukin-10, TGF $\beta$  monoclonal antibody to interleukin-1, ICE inhibitors or depletion via superantigens such as through staphylococcal enterotoxin-A (SEA) induced epidermal Langerhans cell depletion.

Transcutaneous immunization may be induced via the ganglioside GM1 binding activity of CT, LT or subunits such as CTB. Ganglioside GM1 is a ubiquitous cell membrane glycolipid found in all mammalian cells. When the pentameric CT B subunit binds to the cell surface a hydrophilic pore is formed which allows the A subunit to penetrate across the lipid bilayer.

We have shown that transcutaneous immunization by CT or CTB may require ganglioside GM1 binding activity. When mice are transcutaneously immunized with CT, CTA and CTB, only CT and CTB resulted in an immune response. CTA contains the ADP-ribosylating exotoxin activity but only CT and CTB containing the binding activity are able to induce an immune response indicating that the B subunit was necessary and sufficient to immunize through the skin. We conclude that the Langerhans cell or other immune cells may be activated by CTB binding to its cell surface, but more activated by the concurrent presence of the A-subunit.

In addition to activation of the skin component in the immunization process of the present invention, the antigen and/or adjuvant may be activated to enhance immunization. When CT is secreted, cleavage occurs at the trypsin recognition site and the toxin is activated. LT however, is secreted with its trypsin recognition site intact. When LT is secreted in the gastrointestinal tract and thereby exposed to gastrointestinal agents such as trypsin, the proteolytically sensitive residues that join A1 and A2 subunits of LT are cleaved, allowing the A1 subunit to ADP-ribosylate G proteins and therefore exert its toxic effects. The lack of trypsin or related agents in the skin may prevent trypsin cleavage of the proteolytically sensitive residues that join A1 and A2 subunits of LT, diminishing its adjuvant activity.

These two bacterial enterotoxins have many features in common. LT and CT have the same subunit number (A2:B5) and arrangement, and the same biological mechanism of action. An amino acid sequence similarity of 75-77% is found for both chains when comparing LT and CT, and the most significant difference occurs in the respective A chains at positions 192-195. At this site the cleavage of the A chain by trypsin occurs and the site is situated between two cysteine residues that form the internal disulfide bond of the A chain. See, for example, Mekalanos et al. (1979), Spangler (1992), and Sniderman (1995). We propose that these structural differences between the molecules are a significant influence not only on their enterotoxic properties, but also on their ability to function as adjuvants.

Unlike CT produced by *V. cholerae*, LT is not fully biologically active when first isolated from the bacterial cell. Consistent with the A-B model for bacterial toxins, LT requires trypsin proteolysis and disulfide reduction to be fully active (Sniderman, 1995). In the absence of proteolytic processing, the enzymatically active A1 moiety is unable to dissociate from the A2 component and cannot reach its target substrate (adenylate cyclase) on the basolateral surface of the intestinal epithelial cell. This difference in activation of the isolated material results in differences in response thresholds for LT and CT in biologic systems. For instance, CT induces detectable net fluid secretion in the mouse intestine at a dose of 5 to 10 µg. LT induces detectable net secretion in this assay at 50 to 100 µg. In the rabbit ligated ileal loop, the difference is more dramatic and clear cut. Significantly however, when LT is exposed to proteolytic enzymes with trypsin-like specificity, the molecule becomes indistinguishable from CT in any biologic assay system (Clements and Finkelstein, 1979; Dickenson and Clements, 1995).

According to Spangler (1992, citations omitted):

“Subunit A is synthesized as a single polypeptide in both *V. cholerae* and *E. coli*. CTA is proteolytically “nicked” between residues 192 and 195 during secretion from the vibrio by *V. cholerae* hemagglutinin/protease, giving rise to two polypeptides, A1 (Mr = 28,826) and A2 (Mr = 5,407), covalently linked through a disulfide bridge between residues 187 and 199. In contrast, LT remains in the *E. coli* periplasm and is not nicked. Introduced into a genetically engineered strain of *V. cholerae*, LT remained unnicked, although it was secreted in the same manner as CT. Proteolytic processing is therefore not a prerequisite for secretion. Purified LTh can, however, be nicked in vitro, suggesting that the mutant vibrio used by Hirst et al. contained insufficient soluble hemagglutinin to catalyze nicking, rather than indicating an inability of LTA to be nicked. CT, when introduced via an engineered plasmid into *E. coli*, remains unnicked and cell associated in *E. coli*. Therefore, the defect in processing of CT and LT in *E. coli* is related to the failure of *E. coli* to nick and secrete either toxin. This defect may explain the reduced severity of *E. coli*-induced enteric disease when compared with cholera. In both CT and LT, the disulphide bond linking A1 to A2 remains unreduced and the toxin is therefore essentially inactive, until it enters a cell.

“Both the intact A subunit and the holotoxin are relatively inactive ADP-ribosyltransferases compared with the A1 polypeptide. Catalytic activity requires the reduction of the disulfide bond (A1:Cys-187-A2:Cys-199) linking A1 to A2. The cleavage (nicking) between residues A1-Arg 192 and the start of the A2 polypeptide at A2:Met-195 takes place during secretion of CT from the vibrio. tryptic

digestion serves the purpose in vitro for LT. Reduction, which releases CTA1 from CTA2, may be accomplished by a variety of agents, usually dithiothreitol or 2-mercaptoethanol invitro or a thiol:protein oxireductase. The endogenous reducing agent and mechanisms of reduction are not known. An observed time lag of about 16 min between the apparent binding of the toxin to the membrane receptor and the first appearance of the modified substrate intracellularly may be related to the time required for this step to occur following or during insertion or translocation”.

- 10 LTh stands for LT holoezyme. Thus, if trypsin-treated LT were to be used for transcutaneous immunization, we propose similar mechanisms for disrupting disulphide bonds would occur. This may be shown for trypsin activation of LT in which trypsin-activated LT is similarly potent or of greater potency compared to CT and much greater in potency to untreated LT in the mouse Y-1 bioassay (see Dickinson and Clements, 1995).

We propose to activate components of the formulation such as LT using trypsin or similar compounds prior to application the skin to enhance the adjuvant activity and immunogenicity of LT. Activation of LT could also be expected to enhance the immune response to LT as an antigen. The activated adjuvant for transcutaneous immunization is preferably an ADP-ribosylating exotoxin. Optionally, hydration or occlusive dressings may be used in the transcutaneous delivery system in addition to the activation of the adjuvant.

- In addition, LT has an unusual affinity for carbohydrate-containing matrices. Specifically, LT binds to an array of biological molecules containing galactose, including glycoproteins and lipopolysaccharides. This lectin-like binding property of LT results in a broader receptor distribution on mammalian cells for LT than for CT, which binds only to GM1. The two molecules also have many immunologic differences, as demonstrated by immunodiffusion studies, against LT-associated E. coli diarrhea in volunteers receiving B-subunit whole whole-cell cholera vaccine. LT and CT induce different helper T-cell responses. When used as a mucosal adjuvant, CT selectively induces in some cases Th2-type cells in Peyer patches and spleens as manifested by production of interleukins 4 and 5, but not interleukin 2 or gamma interferon; while LT induces both Th1 and Th2 cells and predominantly antigen-specific IgA responses. Taken together, these findings demonstrate that LT and CT are unique molecules, despite their apparent structural similarities. Such differential

behavior makes the ability to activate LT so that it has potency similar to CT useful in manipulating the type of immune response produced to both the toxin itself and to antigens for which LT can be used as an adjuvant. It may also be possible that genetically altered toxoids such as mutants of the trypsin cleavage site may be active by transcutaneous immunization. Such a mutant toxin may be useful as it avoids the risks associated with ingestion or inhaling native toxins.

In a similar manner, PT may be activated to enhance its adjuvant and antigen activities. The S1 subunit of the hexameric PT protein contains the ADP-ribosyltransferase activity while the remaining subunits constitute the B domain.

Similar to LT, PT has both trypsin cleavage sites and disulphide binding sites that play a role in association of the S1 subunit with the B oligomer. It is conceivable that activation by trypsin cleavage, disruption of the disulphide bond or both may enhance the adjuvant and antigen activities of PT in the context of transcutaneous immunization. Activation may also take the form of targeting, achieved by disruption of the hexamer into subunits. For example, the PT subunit S3 binds exclusively to the glycolipids of monocytes and could be used to target Langerhans cells in the skin.

Activation of the antigen or adjuvant could be extended to the concept of transcutaneous immunization using DNA by production of a fusion protein comprised of antigen and adjuvant domains. By this method a plasmid encoding an ADP-ribosylating exotoxin such as CT or LT and constructed to express a separate antigen such as a malaria or HIV antigen simultaneously could be placed on the skin in a hydrating solution or occlusive patch, and then taken up by Langerhans cells.

Expression of the an ADP-ribosylating exotoxin component of the fusions protein such as CT or LT could activate the Langerhans cell, causing it to migrate and present antigen in the lymph node and thereby induce an immune response to the encoded antigen. Another embodiment could include the conjugation of an adjuvant with a plasmid; an Fc portion of IgG to a plasmid to target APCs. A similar immunization could be achieved using separate plasmids for expressing an ADP-ribosylating exotoxin such as CT or LT and another for expressing the antigen such as a malaria or HIV antigen. It is conceivable that multiple genes on a single construct for multiple antigens could be used or multiple plasmids could be used to simultaneously deliver antigens for multivalent immunization. Plasmids encoding other molecules or compounds such as chemokines (e.g., defensins 1 or 2, RANTES, MIP1- $\alpha$ , MIP-2,

interleukin-8) or a cytokine (e.g., interleukin-1 $\beta$ , -2, -6, -10 or -12;  $\gamma$ -interferon; tumor necrosis factor- $\alpha$ ; or granulocyte-monocyte-colony stimulating factor) (reviewed in Nohria and Rubin, 1994), a heat shock protein or a derivative, a derivative of *Leishmania major* LeIF (Skeiky et al., 1995), cholera toxin toxin B, a  
5 lipopolysaccharide (LPS) derivative (e.g., lipid A or monophosphoryl lipid A), or superantigen (Saloga et al., 1996), or other ADP-ribosylating exotoxins might be delivered with protein antigens.

Other means of activating the transcutaneous adjuvants may be effective, such as adding detergents and phospholipid to the formulation to enhance CT activity by  
10 ADP-ribosylation factor (see, for example, Spangler, 1992).

For immunization using adjuvant or antigen activation, modification of the adjuvant or antigen component of the formulation may reduce its effectiveness in parenteral immunization without destroying the utility of the formulation in transcutaneous immunization when the adjuvant and/or antigen is activated.  
15 Undesirable properties (e.g., toxicity, allergic reactivity, other side effects) of the adjuvant or antigen in the formulation may be reduced by modification without destroying its effectiveness in transcutaneous immunization. Activation of such modified adjuvant or antigen may involve, for example, removal of a reversible chemical modification (e.g., proteolysis) or a coating which reversibly isolates a  
20 component of the formulation from the immune system (i.e., an encapsulated formulation). Alternatively, the adjuvant and/or antigen comprising the formulation may be encapsulated in a particle (e.g., microspheres, nanoparticles). Phagocytosis of a particle may, by itself, enhance activation of an antigen presenting cell by upregulating expression of major histocompatibility antigens and/or costimulatory  
25 molecules (e.g., MHC class II, B7-2).



## ANTIGEN

Antigen of the invention may be expressed by recombinant means, preferably as a fusion with an affinity or epitope tag (Summers and Smith, 1987; Goeddel, 1990; Ausubel et al., 1996); chemical synthesis of an oligopeptide, either free or conjugated to carrier proteins, may be used to obtain antigen of the invention (Bodanszky, 1993; Wisdom, 1994). Oligopeptides are considered a type of polypeptide. Oligopeptide lengths of 6 residues to 20 residues are preferred. Polypeptides may also be synthesized as branched structures such as those disclosed in U.S. Pat. Nos. 5,229,490 and 5,390,111. Antigenic polypeptides include, for example, synthetic or recombinant B-cell and T-cell epitopes, universal T-cell epitopes, and mixed T-cell epitopes from one organism or disease and B-cell epitopes from another. Antigen obtained through recombinant means or peptide synthesis, as well as antigen of the invention obtained from natural sources or extracts, may be purified by means of the antigen's physical and chemical characteristics, preferably by fractionation or chromatography (Janson and Ryden, 1989; Deutscher, 1990; Scopes, 1993). A multivalent antigen formulation may be used to induce an immune response to more than one antigen at the same time. Conjugates may be used to induce an immune response to multiple antigens, to boost the immune response, or both. Additionally, toxins may be boosted by the use of toxoids, or toxoids boosted by the use of toxins. Transcutaneous immunization may be used to boost responses induced initially by other routes of immunization such as by oral, nasal or parenteral routes. Antigen includes, for example, toxins, toxoids, subunits thereof, or combinations thereof (e.g., cholera toxin, tetanus toxoid); additionally, toxins, toxoids, subunits thereof, or combinations thereof may act as both antigen and adjuvant.

Antigen may be solubilized in a buffer. Suitable buffers include, but are not limited to, phosphate buffered saline  $\text{Ca}^{++}/\text{Mg}^{++}$  free (PBS), normal saline (150 mM NaCl in water), and Tris buffer. Glycerol may be a suitable non-aqueous buffer for use in the present invention. Antigen may also be in suspension. The detergent may be left in the immunizing solution to enhance penetration.

Hydrophobic antigen can be solubilized in a detergent, for example a polypeptide containing a membrane-spanning domain. Furthermore, for formulations containing liposomes, an antigen in a detergent solution (e.g., a cell membrane

extract) may be mixed with lipids, and liposomes then may be formed by removal of the detergent by dilution, dialysis, or column chromatography. See, Gregoriadis (1993). Certain antigens such as, for example, those from a virus (e.g., hepatitis A) need not be soluble per se, but can be incorporated directly into a lipid membrane (e.g., a virosome as described by Morein and Simons, 1985), in a suspension of virion alone, or suspensions of microspheres nanoparticles or heat-inactivated bacteria which may be taken up by and activate antigen presenting cells (e.g., opsonization).

Plotkin and Mortimer (1994) provide antigens which can be used to vaccinate animals or humans to induce an immune response specific for particular pathogens, as well as methods of preparing antigen, determining a suitable dose of antigen, assaying for induction of an immune response, and treating infection by a pathogen (e.g., bacterium, virus, fungus, or parasite).

Bacteria include, for example: anthrax, campylobacter, cholera, clostridia, diphtheria, enterotoxigenic *E. coli*, giardia, gonococcus, *Helicobacter pylori* or urease produced by *H. pylori* (Lee and Chen, 1994), *Hemophilus influenza B*, *Hemophilus influenza non-typable*, meningococcus, mycobacterium, pertussis, pneumococcus, salmonella, shigella, staphylococcus, *Streptococcus B*, tetanus, *Vibrio cholerae*, *Borrelia burgdorfi* and *Yersinia*; and products thereof.

Viruses include, for example: adenovirus, dengue serotypes 1 to 4 (Delenda et al., 1994; Fonseca et al., 1994; Smucny et al., 1995), ebola (Jahrling et al., 1996), enterovirus, hanta virus, hepatitis serotypes A to E (Blum, 1995; Katkov, 1996; Lieberman and Greenberg, 1996; Mast, 1996; Shafara et al., 1995; Smedila et al., 1994; U.S. Pat. Nos. 5,314,808 and 5,436,126), herpes simplex virus 1 or 2, human immunodeficiency virus (Deprez et al., 1996), human papilloma virus, influenza, measles, Norwalk, Japanese equine encephalitis, papilloma virus, parvovirus B19, polio, rabies, respiratory syncytial virus, rotavirus, rubella, rubeola, St. Louis encephalitis, vaccinia, vaccinia constructs containing genes coding for other antigens such as malaria antigens, varicella, and yellow fever; and products thereof.

Parasites include, for example: *Entamoeba histolytica* (Zhang et al., 1995); *Plasmodium* (Bathurst et al., 1993; Chang et al., 1989, 1992, 1994; Fries et al., 1992a, 1992b; Herrington et al., 1991; Khusmith et al., 1991; Malik et al., 1991; Migliorini et al., 1993; Pessi et al., 1991; Tam, 1988; Vreden et al., 1991; White et al., 1993;

Wiesmueller et al., 1991), Leishmania (Frankenburg et al., 1996), and the Helminthes; and products thereof.

Other viruses which can be used in the present invention are disclosed in Gordon, 1997 and include, for example, Adenovirus (respiratory disease),

5 Coronavirus (respiratory and enteric disease), Cytomegalovirus (mononucleosis), Dengue virus (dengue fever, shock syndrome), Epstein-Barr virus (mononucleosis, Burkitt's lymphoma), Hepatitis A, B and C virus (liver disease), Herpes simplex virus type 1 (encephalitis, stomatitis), Herpes simplex virus type 2 (genital lesions), Human herpesvirus-6 (unknown, possibly Kaposi's sarcoma), Human immunodeficiency

10 virus types 1 and 2 (acquired immunodeficiency syndrome-AIDS), Human T-cell lymphotropic virus type 1 (T-cell leukemia), Influenza A, B, and C (respiratory disease), Japanese encephalitis virus (pneumonia, encephalopathy), Measles virus (subacute sclerosing panencephalitis), Mumps virus (meningitis, encephalitis), Papillomavirus (warts, cervical carcinoma), Parvovirus (respiratory disease, anemia),

15 Poliovirus (paralysis), Polyomavirus JC (multifocal leukoencephalopathy), Polyomavirus BK (hemorrhagic cystitis), Rabies virus (nerve dysfunction), Respiratory syncytial virus (respiratory disease), Rhinovirus (common cold), Rotavirus (diarrhea), Rubella virus (fetal malformations), Vaccinia virus (generalized infection), Yellow fever virus (jaundice, renal and hepatic failure), Varicella zoster

20 virus (chickenpox).

Other bacteria which can be used in the present invention are disclosed in Gordon, 1997 and include, for example, Bacillus anthracis (anthrax), Bordetella pertussis (whooping cough), Borrelia burgdorferi (lyme disease), Campylobacter jejuni (gastroenteritis), Chlamydia trachomatis (pelvic inflammatory disease,

25 blindness), Clostridium botulinum (botulism), Corynebacterium diphtheriae (diphtheria), Escherichia coli (diarrhea, urinary tract infections), Haemophilus influenzae (pneumonia), Helicobacter pylori (gastritis, duodenal ulcer), Legionella pneumophila (Legionnaires' disease), Listeria monocytogenes (meningitis, sepsis), Mycobacterium leprae (leprosy), Mycobacterium tuberculosis (tuberculosis),

30 Neisseria gonorrhoeae (gonorrhea), Neisseria meningitidis (sepsis, meningitis), Pseudomonas aeruginosa (nosocomial infections), Pseudomonas aeruginosa (nosocomial infections), Rickettsia (Rocky Mountain spotted fever), Salmonella (typhoid fever, gastroenteritis), Shigella (dysentery), Staphylococcus aureus

(impetigo, toxic shock syndrome), *Streptococcus pneumoniae* (pneumonia, otitis media), *Streptococcus pyogenes* (Rheumatic fever, pharyngitis), *Treponema pallidum* (syphilis), *Vibrio cholerae* (cholera), *Yersinia pestis* (bubonic plague).

Other parasites which can be used in the present invention are disclosed in  
5 Gordon, 1997 and include, for example, African trypanosomes (trypanosomiasis),  
Entamoeba histolytica (amebic dysentery), Giardia lamblia (diarrheal disease),  
Leishmania (lesions of the spleen, tropical sores), Plasmodium (malaria),  
Microfilariae (filariasis), Schistosomes (schistosomiasis), Toxoplasma gondii  
(toxoplasmosis), Trichomonas vaginalis (vaginitis), Trypanosoma cruzi (Chagas  
10 disease).

Fungi which can be used in the present invention are disclosed in Gordon,  
1997 and include, for example, Candida albicans (mucosal infections), Histoplasma  
(lung, lymph node infections), Pneumocystis carinii (pneumonia in AIDS),  
Aspergillus fumigatis (aspergillosis).

#### ADJUVANT

The formulation also contains an adjuvant, although a single molecule may  
contain both adjuvant and antigen properties (e.g., cholera toxin) (Elson and  
Dertzbaugh, 1994). Adjuvants are substances that are used to specifically or non-  
20 specifically potentiate an antigen-specific immune response. Usually, the adjuvant  
and the formulation are mixed prior to presentation of the antigen but, alternatively,  
they may be separately presented within a short interval of time.

Adjuvants are usually products of bacteria, parasites or even viruses or  
bacteria, but may be derived from other natural or synthetic sources. Adjuvants  
25 include, for example, an oil emulsion (e.g., complete or incomplete Freund's  
adjuvant), a chemokine (e.g., defensins 1 or 2, RANTES, MIP1- $\alpha$ , MIP-2, interleukin-  
8) or a cytokine (e.g., interleukin-1 $\beta$ , -2, -6, -10 or -12;  $\gamma$ -interferon; tumor necrosis  
factor- $\alpha$ ; or granulocyte-monocyte-colony stimulating factor) (reviewed in Nohria and  
Rubin, 1994), a muramyl dipeptide derivative (e.g., murabutide, threonyl-MDP or  
30 muramyl tripeptide), a heat shock protein or a derivative, a derivative of Leishmania  
major LeIF (Skeiky et al., 1995), cholera toxin or cholera toxin B, bARES, a  
lipopolysaccharide (LPS) derivative (e.g., lipid A or monophosphoryl lipid A ,  
synthetic lipid A analogues), or superantigen (Saloga et al., 1996) block copolymers or

other polymers known in the art. Also, see Richards et al. (1995) for other adjuvants useful in immunization.

An adjuvant may be chosen to preferentially induce antibody or cellular effectors, specific antibody isotypes (e.g., IgM, IgD, IgA1, IgA2, secretory IgA, IgE, IgG1, IgG2, IgG3, and/or IgG4), or specific T-cell subsets (e.g., CTL, Th1, Th2 and/or T<sub>DTH</sub>) (Munoz et al., 1990; Glenn et al., 1995).

CpGs are among a class of structures which have patterns allowing the immune system to recognize their pathogenic origins to stimulate the innate immune response leading to adaptive immune responses (Medzhitov and Janeway, 1997). These structures are called pathogen-associated molecular patterns (PAMPs) and include lipopolysaccharides, teichoic acids, unmethylated CpG motifs, double stranded RNA and mannins, for example.

PAMPs induce endogenous danger signals that can enhance the immune response, act as costimulators of T-cell function and control the effector function. The ability of PAMPs to induce these responses play a role in their potential as adjuvants and their targets are APCs such as macrophages and dendritic cells. The antigen presenting cells of the skin could likewise be stimulated by PAMPs transmitted through the skin. For example, Langerhans cells, a type of dendritic cell, could be activated by a PAMP in solution on the skin with a transcutaneously poorly immunogenic molecule and be induced to migrate and present this poorly immunogenic molecule to T-cells in the lymph node, inducing an antibody response to the poorly immunogenic molecule. PAMPs could also be used in conjunction with other skin adjuvants such as cholera <sup>toxin</sup> ~~toxin~~ to induce different costimulatory molecules and control different effector functions to guide the immune response, for example from a Th2 to a Th1 response.

Cholera toxin is a bacterial exotoxin from the family of ADP-ribosylating exotoxins (referred to as bAREs). Most bAREs are organized as A:B dimer with a binding B subunit and an A subunit containing the ADP-ribosyltransferase. Such toxins include diphtheria, Pseudomonas exotoxin A, cholera toxin (CT), E. coli heat-labile enterotoxin (LT), pertussis toxin (PT), C. botulinum toxin C2, C. botulinum toxin C3, C. limosum exoenzyme, B. cereus exoenzyme, Pseudomonas exotoxin S, Staphylococcus aureus EDIN, and B. sphaericus toxin.

Cholera toxin is an example of a bARE that is organized with A and B subunits. The B subunit is the binding subunit and consists of a B-subunit pentamer which is non-covalently bound to the A subunit. The B-subunit pentamer is arranged in a symmetrical doughnut-shaped structure that binds to GM1-ganglioside on the target cell. The A subunit serves to ADP ribosylate the alpha subunit of a subset of the hetero trimeric GTP proteins (G proteins) including the Gs protein which results in the elevated intracellular levels of cyclic AMP. This stimulates release of ions and fluid from intestinal cells in the case of cholera.

Cholera toxin (CT) and its B subunit (CTB) have adjuvant properties when used as either an intramuscular or oral immunogen (Elson and Dertzbaugh, 1994; Trach et al., 1997). Heat-labile enterotoxin from *E. coli* (LT) is 75-77% homologous at the amino acid level with CT and possesses similar binding properties; it also appears to bind the GM1-ganglioside receptor in the gut and has similar ADP-ribosylating exotoxin activities. Another bARE, *Pseudomonas* exotoxin A (ETA), binds to the  $\alpha$ 2-macroglobulin receptor-low density lipoprotein receptor-related protein (Kounnas et al., 1992). bAREs are reviewed by Krueger and Barbieri (1995). CT, CTB, LT, ETA and PT, despite having different cellular binding sites, are potent adjuvants for transcutaneous immunization, inducing high levels of IgG antibodies but not IgE antibodies. CTB without CT can also induce high levels of IgG antibodies. Thus, both bAREs and a derivative thereof can effectively immunize when epicutaneously applied to the skin in a simple solution.

All licensed vaccines require a level of antibody for approval – no other immune component such as T-cell proliferation is used. Protection against the life-threatening infections diphtheria, pertussis, and tetanus (DPT) can be achieved by inducing high levels of circulating anti-toxin antibodies. Pertussis may be an exception in that some investigators feel that antibodies directed to other portions of the invading organism are necessary for protection, although this is controversial (see Schneerson et al., 1996) and most new generation acellular pertussis vaccines have PT as a component of the vaccine (Krueger and Barbieri, 1995). The pathologies in the diseases caused by DPT are directly related to the effects of their toxins and anti-toxin antibodies most certainly play a role in protection (Schneerson et al., 1996).

In general, toxins can be chemically inactivated to form toxoids which are less toxic but remain immunogenic. We envision that the one embodiment of

transcutaneous immunization system will use toxin-based immunogens and adjuvants to achieve anti-toxin levels adequate for protection against these diseases. The anti-toxin antibodies may be induced through immunization with the toxins, or genetically-detoxified toxoids themselves, or with toxoids and adjuvants such as CT. Genetically

5 toxoided toxins which have altered ADP-ribosylating exotoxin activity, or trypsin cleavage site mutations or other mutations are envisioned to be especially useful as non-toxic activators of antigen presenting cells used in transcutaneous immunization. Mutants based on inactivating the catalytic activity of the ADP-ribosyl transferase by genetic deletion retain the binding capabilities, but lack the toxicity, of the natural

10 toxins. This approach is described by Burnette et al. (1994), Rappuoli et al. (1995), and Rappuoli et al. (1996). Such genetically toxoided exotoxins could be useful for transcutaneous immunization system in that they would not create a safety concern as the toxoids would not be considered toxic. There are other genetically altered toxins which have, for example, deletions of the trypsin cleavage site and use both non-toxic

15 and immunogenic on the skin. However, activation through a technique such as trypsin cleavage would be expected to enhance the adjuvant qualities of LT through the skin which is lacking inherent trypsin enzymes. Additionally, several techniques exist to chemically toxoid toxins which can address the same problem (Schneerson et al., 1996). These techniques could be important for certain applications, especially

20 pediatric applications, in which ingested toxins (e.g., diphtheria toxin) might possibly create adverse reactions.

Optionally, an activator of Langerhans cells may be used as an adjuvant. Examples of such activators include: an inducer of heat shock protein; contact sensitizer (e.g., trinitrochlorobenzene, dinitrofluorobenzene, nitrogen mustard,

25 pentadecylcatechol); toxin (e.g., Shiga toxin, Staph enterotoxin B); lipopolysaccharide, lipid A, or derivatives thereof; bacterial DNA (Stacey et al., 1996); cytokine (e.g., tumor necrosis factor- $\alpha$ , interleukin-1 $\beta$ , -10, -12); calcium ions in solution; calcium ionophores, and chemokine (e.g., defensins 1 or 2, RANTES, MIP-1 $\alpha$ , MIP-2, interleukin-8).

30 If an immunizing antigen has sufficient Langerhans cell activating capabilities then a separate adjuvant may not be required, as in the case of CT which is both antigen and adjuvant. It is envisioned that whole cell preparations, live viruses, attenuated viruses, DNA plasmids, and bacterial DNA could be sufficient to

immunize transcutaneously then an adjuvant is present. It may be possible to use low concentrations of contact sensitizers or other activators of Langerhans cells to induce an immune response without inducing skin lesions.

## 5 PRACTICAL ASPECTS OF TRANSCUTANEOUS IMMUNIZATION

Efficient immunization can be achieved with the present invention because transcutaneous delivery of antigen may target the Langerhans cell. These cells are found in abundance in the skin and are efficient antigen presenting cells leading to T-cell memory and potent immune responses. Because of the presence of large numbers of Langerhans cells in the skin, the efficiency of transcutaneous delivery may be related to the surface area exposed to antigen and adjuvant. In fact, the reason that transcutaneous immunization is so efficient may be that it targets a larger number of these efficient antigen presenting cells than intramuscular immunization.

15 We envision the present invention will enhance access to immunization, while inducing a potent immune response. Because transcutaneous immunization does not involve physical penetration of the skin and the complications and difficulties thereof, the requirements of trained personnel, sterile technique, and sterile equipment are reduced. Furthermore, the barriers to immunization at multiple sites or to multiple immunizations are diminished. Immunization by a single application of the formulation is also envisioned, but boosting is generally needed. Needle free immunization is a priority for the World Health Organization (WHO) because of the reuse of needles which causes needle-borne disease.

25 Immunization may be achieved using epicutaneous application of a simple solution of antigen and adjuvant impregnated in gauze under an occlusive patch, or by using other patch technologies; creams, gels, immersion, ointments and sprays are other possible methods of application. The immunization could be given by untrained personnel, and is amenable to self-application. Large-scale field immunization could occur given the easy accessibility to immunization. Additionally, a simple immunization procedure would improve access to immunization by pediatric patients and the elderly, and populations in Third World countries.

For previous vaccines, their formulations were injected through the skin with needles. Injection of vaccines using needles carries certain drawbacks including the



need for sterile needles and syringes, trained medical personnel to administer the vaccine, discomfort from the injection, and potential complications brought about by puncturing the skin with the needle. Immunization through the skin without the use of needles (i.e., transcutaneous immunization) represents a major advance for vaccine delivery by avoiding the aforementioned drawbacks.

Moreover, transcutaneous immunization may be superior to immunization using needles as more immune cells would be targeted by the use of several locations targeting large surface areas of skin. A therapeutically effective amount of antigen sufficient to induce an immune response may be delivered transcutaneously either at a single cutaneous location, or over an area of intact skin covering multiple draining lymph node fields (e.g., cervical, axillary, inguinal, epitrocheal, popliteal, those of the abdomen and thorax). Such locations close to numerous different lymphatic nodes at locations all over the body will provide a more widespread stimulus to the immune system than when a small amount of antigen is injected at a single location by intradermal subcutaneous or intramuscular injection.

Antigen passing through or into the skin may encounter antigen presenting cells which process the antigen in a way that induces an immune response. Multiple immunization sites may recruit a greater number of antigen presenting cells and the larger population of antigen presenting cells that were recruited would result in greater induction of the immune response. It is conceivable that absorption through the skin may deliver antigen to phagocytic cells of the skin such as, for example, dermal dendritic cells, macrophages, and other skin antigen presenting cells; antigen may also be delivered to phagocytic cells of the liver, spleen, and bone marrow that are known to serve as the antigen presenting cells through the blood stream or lymphatic system. Langerhans cells, dendritic cells, and macrophages may be specifically targeted using Fc receptor conjugated to or recombinantly produced as a protein fusion with adjuvant; also, complement receptors (C3, C5) may be conjugated to or recombinantly produced as a protein fusion with protein A or protein G to target surface immunoglobulin of B cells. The result would be targeted distribution of antigen to antigen presenting cells to a degree that is rarely, if ever achieved, by current immunization practices.

The transcutaneous immunization system may be applied directly to the skin and allowed to air dry; rubbed into the skin or scalp; held in place with a dressing,

patch, or absorbent material; immersion; otherwise held by a device such as a stocking, slipper, glove, or shirt; or sprayed onto the skin to maximize contact with the skin. The formulation may be applied in an absorbant dressing or gauze. The formulation may be covered with an occlusive dressing such as, for example,

5 AQUAPHOR (an emulsion of petrolatum, mineral oil, mineral wax, wool wax, panthenol, bisabol, and glycerin from Beiersdorf, Inc.), plastic film, COMFEEL (Coloplast) or vaseline; or a non-occlusive dressing such as, for example, DUODERM (3M) or OPSITE (Smith & Napheu). An occlusive dressing completely excludes the passage of water. The formulation may be applied to single or multiple sites, to single

10 or multiple limbs, or to large surface areas of the skin by complete immersion. The formulation may be applied directly to the skin.

Genetic immunization has been described in U.S. Pat. Nos. 5,589,466, 5,593,972, and 5,703,055. The nucleic acid(s) contained in the formulation may encode the antigen, the adjuvant, or both. It would generally be expected that the

15 immune response would be enhanced by the coadministration of an adjuvant, for example, CT, LT or CpGs to the nucleic acid encoding for the antigen. The nucleic acid may or may not be capable of replication; it may be non-integrating and non-infectious. For example, the nucleic acid may encode a fusion polypeptide comprising antigen and a ubiquitin domain to direct the immune response to a class I restricted

20 response. The nucleic acid may further comprise a regulatory region (e.g., promoter, enhancer, silencer, transcription initiation and termination sites, RNA splice acceptor and donor sites, polyadenylation signal, internal ribosome binding site, translation initiation and termination sites) operably linked to the sequence encoding the antigen. The nucleic acid may be complexed with an agent that promotes transfection such as

25 cationic lipid, calcium phosphate, DEAE-dextran, polybrene-DMSO, or a combination thereof; also, immune cells can be targeted by conjugation of DNA to Fc receptor or protein A/G, or encapsulating DNA in an agent linked to Fc receptor or protein A/G. The nucleic acid may comprise regions derived from viral genomes. Such materials and techniques are described by Kriegler (1990) and Murray (1991).

30 An immune response may comprise humoral (i.e., antigen-specific antibody) and/or cellular (i.e., antigen-specific lymphocytes such as B cells, CD4<sup>+</sup> T cells, CD8<sup>+</sup> T cells, CTL, Th1 cells, Th2 cells, and/or T<sub>DTH</sub> cells) effector arms. Moreover, the

immune response may comprise NK cells that mediate antibody-dependent cell-mediated cytotoxicity (ADCC).

The immune response induced by the formulation of the invention may include the elicitation of antigen-specific antibodies and/or cytotoxic lymphocytes (CTL, reviewed in Alving and Wassef, 1994). Antibody can be detected by immunoassay techniques, and the detection of various isotypes (e.g., IgM, IgD, IgA1, IgA2, secretory IgA, IgE, IgG1, IgG2, IgG3, or IgG4) may be expected. An immune response can also be detected by a neutralizing assay. Antibodies are protective proteins produced by B lymphocytes. They are highly specific, generally targeting one epitope of an antigen. Often, antibodies play a role in protection against disease by specifically reacting with antigens derived from the pathogens causing the disease.

CTLs are particular protective immune cells produced to protect against infection by a pathogen. They are also highly specific. Immunization may induce CTLs specific for the antigen, such as a synthetic oligopeptide based on a malaria protein, in association with self-major histocompatibility antigen. CTLs induced by immunization with the transcutaneous delivery system may kill pathogen infected cells. Immunization may also produce a memory response as indicated by boosting responses in antibodies and CTLs, lymphocyte proliferation by culture of lymphocytes stimulated with the antigen, and delayed type hypersensitivity responses to intradermal skin challenge of the antigen alone.

In a viral neutralization assay, serial dilutions of sera are added to host cells which are then observed for infection after challenge with infectious virus. Alternatively, serial dilutions of sera may be incubated with infectious titers of virus prior to inoculation of an animal, and the inoculated animals are then observed for signs of infection.

The transcutaneous immunization system of the invention may be evaluated using challenge models in either animals or humans, which evaluate the ability of immunization with the antigen to protect the subject from disease. Such protection would demonstrate an antigen-specific immune response. In lieu of challenge, for example achieving anti-diphtheria antibody titers of 5 IU/ml or greater is generally assumed to indicate optimum protection and serves as a surrogate marker for protection (Plotkin and Mortimer, 1994).

Vaccination has also been used as a treatment for cancer and autoimmune disease. For example, vaccination with a tumor antigen (e.g., prostate specific antigen) may induce an immune response in the form of antibodies, CTLs and lymphocyte proliferation which allows the body's immune system to recognize and kill tumor cells. Tumor antigens useful for vaccination have been described for melanoma (U.S. Pat. Nos. 5,102,663, 5,141,742, and 5,262,177), prostate carcinoma (U.S. Pat. No. 5,538,866), and lymphoma (U.S. Pat. Nos. 4,816,249, 5,068,177, and 5,227,159). Vaccination with T-cell receptor oligopeptide may induce an immune response that halts progression of autoimmune disease (U.S. Pat. Nos. 5,612,035 and 5,614,192; Antel et al., 1996; Vandenbark et al., 1996). U.S. Pat. No. 5,552,300 also describes antigens suitable for treating autoimmune disease.

The following is meant to be illustrative of the present invention; however, the practice of the invention is not limited or restricted in any way by the examples.

## EXAMPLES

**Immunization procedure.** Twenty four hours prior to immunization, the back of the mouse is shaved from the distal aspect of the scapula to 0.5 cm above the base of the tail. In the case of C57BL/6 mice, the animals are lightly anesthetized (40 mg/kg ketamine : 4 mg/kg xylazine mixture in saline) prior to shaving. On the day of immunization the animals are immunized with .04 ml of an anesthesia mixture (2.3 mL sterile saline (Sigma): 5 mL ketamine (100 mg/mL, Parke-Davis) : 0.5 mL xylazine (100 mg/mL, Phoenix Pharmaceuticals)) which delivers a final dose of approximately 110 mg/kg ketamine and 11 mg/kg xylazine. For procedures requiring alcohol swabbing, the back is wiped 10x (5x sweeping up the back towards the head, flip over alcohol pad and sweep back 5x more) using an isopropyl pad. The alcohol is allowed to evaporate for 5 minutes. Hydration of the back is accomplished by gently rubbing the back with a sterile water-saturated gauze pad so as to form a pool of water on the back. After a 5 minute hydration period, the back is blotted dry with a dry gauze pad. Next, antigen - generally  $\leq 100 \mu\text{g}$  of antigen and adjuvant in 100  $\mu\text{l}$  final volume, is applied to the back using a pipette and tip and left on the skin for 60 to 120 minutes. After the defined immunization period has been reached, any excess solution in the immunized area is blotted with cotton gauze. The animals are then rinsed animals under a slow steady stream of lukewarm tap water for 10 seconds to remove any excess antigen, blotted dry and the rinsing procedure repeated. The cages

are then placed onto the heating pads until they are fully recovered from the anesthesia.

Measurement of Human anti-LT Antibody Titers. Anti-LT IgG titers were

determined as previously described (Svennerholm A-M., Holmgren, J., Black, R.,

5 Levine, M. & Merson, M. Serologic differentiation between antitoxin responses to infection with *Vibrio cholerae* and enterotoxin-producing *Escherichia coli*. *J. Infect. Dis.* **147**, 541-522 (1983). 96 well (Type-Russell) plates were coated overnight with monosialoganglioside- $G_{M1}$  (Sigma, St. Louis, MO) of LT (Sigma) , blocked with 5% dry milk in PBS-0.05% Tween. Responses were detected using goat anti-human

10 IgG( $\gamma$ )-HRP (Kirkegaard and Perry, Gaithersburg, MD., and 2,2'-azino-di[3-ethylbenzthiazoline sulfonate (Kirkegaard and Perry) as substrate and plates were read at 405 nm. Results are reported in ELISA units (EU) which are defined as the inverse dilution of sample which yields an OD of 1.0. Anti-LT IgA was determined in the same manner as anti-LT IgG except that goat anti-human IgA( $\alpha$ )-HRP (Kirkegaard and Perry) was used as secondary antibody and ODs were plotted against a standard IgA curve yielding results expressed in ng/ml. The standard IgA curve and total serum IgA were determined by using unlabeled goat anti-human IgA (Kirkegaard and Perry) followed by blocking as above and then application of serial dilutions of IgA standard

15  
20 Example 1.

Swabbing the skin with a treated or untreated swab is thought to physically and chemically remove a small portion of the stratum corneum and thus enhance skin penetration. Swabs can be made of materials such as, for example, cotton, nylon, rayon and polyethylene. Alcohol swabbing is thought to remove a small portion of the stratum corneum and acts both as a physical means and chemical means of penetration enhancement. In the example above, the enhancement of the immune response to transcutaneous immunization can be seen with this penetration enhancement method. BALB/c mice 6 to 8 weeks of age were anesthetized and shaved as described in the "immunization procedure". Twenty four hours later, the backs of the animals were either wiped with a gauze pad saturated in water "water" or wiped for approximately 10 seconds with an alcohol prep pad containing 70% isopropyl alcohol "isopropanol". The alcohol was allowed to evaporate for approximately 5 minutes. The excess water was removed from the backs of the

“water” group by blotting. All animals were then treated with 20 µg of CT (100 µl of a 0.2 mg/ml solution). Removal of excess antigen was conducted as described in the “immunization procedure.”

5 The anti-CT antibody titers were determined using ELISA as described above for “ELISA IgG (H+L)” 3 weeks after a single immunization. The results are shown in Table 1. While CT was clearly immunogenic in both groups, the group treated with the alcohol prep pads exhibited a geometric mean titer that was 6 fold higher and the individual titers were more consistent than the “water” animals. Thus it appears that chemical and physical disruption of the skin surface with alcohol swabs enhances  
10 delivery of antigen by the transcutaneous route.

Table 1. Enhancement of transcutaneous immunization by chemical penetration enhancement: Anti-CT titers in mice that had the skin treated with an alcohol prep pad before application of the antigen.

Animal #	anti-CT IgG (H+L) ELISA units		
	treatment	prebleed	week 3
7146	water		1275
7147	water		69
7148	water		7420
7149	water		6025
7150	water		388
geometric mean pooled prebleed		7	1088
7161	isopropanol		3100
7162	isopropanol		14797
7163	isopropanol		6670
7164	isopropanol		7426
7165	isopropanol		7024
geometric mean pooled prebleed		7	6928

## Example 2.

To assess whether chemical penetration enhancement alone might augment transcutaneous immunization a detergent was used on the skin. BALB/c mice 6 to 8 weeks of age were anesthetized and shaved as described in the “immunization procedure.” Twenty-four hours later, the backs of the “water” group were wiped with a gauze pad saturated in water and a pool of water was placed on the back.

Approximately 5 minutes later, any excess water was removed and 25  $\mu$ l of CT (50  $\mu$ l of a 0.5 mg/ml solution) was applied to the back. Alternatively, 24 hours after shaving, the backs of the “5% SDS” group were treated by dripping 300  $\mu$ l of 5% SDS (Sodium Dodecyl Sulfate – a 1 to 1 mixture of deionized water and commercial stock of 10% SDS), a detergent, for approximately 12 minutes followed by blotting off any excess SDS with a dry gauze pad. SDS can be applied to the skin in a carrier such as, for example, a pad and then any excess SDS can be removed with a dry gauze pad. Thereafter the animals were hydrated and immunized as per the “water” group.

Removal of excess antigen was conducted as described in the “immunization procedure.”

The anti-CT antibody titers were determined using ELISA as described above for “ELISA IgG (H+L)” 2 weeks after a single immunization. The results are shown in tables 2a and 2b. While CT was clearly immunogenic in both groups, the geometric mean titer in the 5% SDS treated group approximately 2 fold higher and the titers were more consistent among the latter animals as compared with the “water” animals. Thus it appears that chemical disruption of the skin surface with detergent (5% SDS) enhances delivery of antigen by the transcutaneous route.



Table 2a. Enhancement of transcutaneous immunization by chemical penetration enhancement: Anti-CT titers in mice that had the skin treated with detergent (5% SDS) before application of the antigen.

Animal #	anti-CT IgG (H+L) ELISA units		
	treatment	prebleed	week 2
546	water		4629
547	water		3154
548	water		7288
549	water		1719
550	water		11779
geometric mean pooled prebleed		5	3678
596	5% SDS		6945
597	5% SDS		2244
598	5% SDS		8604
599	5% SDS		7093
600	5% SDS		12583
geometric mean pooled prebleed		1	5553

Table 2b. Enhancement of transcutaneous immunization by chemical penetration enhancement: Anti-CT titers in mice that had the skin treated with detergent (5% SDS) before application of the antigen.

Animal #	anti-CT IgG (H+L) ELISA units		
	treatment	prebleed	week 3
546	water		22525
547	water		8939
548	water		11885
549	water		5121
550	water		37770
geometric mean pooled prebleed		11	10521
596	5% SDS		102387
597	5% SDS		6597
598	5% SDS		47245
599	5% SDS		45565
600	5% SDS		38413
geometric mean pooled prebleed		6	34725

Example 3.

Another form of chemical penetration enhancement, a depilatory (such as, for example, calcium hydroxide or the like) is widely used in dermatologic experiments and was shown to enhance transcutaneous immunization. BALB/c mice 6 to 8 weeks of age were anesthetized and shaved as described in the “immunization procedure.” Twenty-four hours later, the backs of the “water” group were wiped with a gauze pad saturated in water and a pool of water was placed on the back. Approximately 5 minutes later, any excess water was removed and 25 µg of CT (50 µl of a 0.5 mg/ml solution) was applied to the back. Alternatively, twenty-four hours after shaving, the backs of the “nair©” group were treated with 100 µl of nair© cream for approximately 12 minutes followed by wiping off of the formulation with a gauze pad saturated in water. Such treatment can continue for from about 0.1 to 30 minutes preferably about 20 minutes and more preferably about 12 minutes. Thereafter the animals were hydrated and immunized as per the “water” group. Removal of excess antigen was conducted as described in the “immunization procedure.”

The anti-CT antibody titers were determined using ELISA as described above for “ELISA IgG (H+L)” 2 weeks after a single immunization. The results are shown in tables 3a and 3b. While CT was clearly immunogenic in both groups, the geometric mean titer in the nair treated group was 3 fold higher and the titers were more consistent among the latter animals as compared with the “water” animals. Thus it appears that chemical disruption of the skin surface with calcium hydroxide, the active ingredient in nair© cream, enhances delivery of antigen by the transcutaneous route.

Table 3a. Enhancement of transcutaneous immunization by chemical penetration enhancement: Anti-CT titers in mice that had the skin treated with calcium hydroxide (nair©) before application of the antigen.

Animal #	anti-CT IgG (H+L) ELISA units		
	treatment	prebleed	week 2
546	water		4629
547	water		3154
548	water		7288
549	water		1719
550	water		11779
geometric mean pooled prebleed		5	3678
581	nair©		17621
582	nair©		12261
583	nair©		7235
584	nair©		7545
585	nair©		5997
geometric mean pooled prebleed		4	10421

Table 3b. Enhancement of transcutaneous immunization by chemical penetration enhancement: Anti-CT titers in mice that had the skin treated with calcium hydroxide (nair©) before application of the antigen.

Animal #	anti-CT IgG (H+L) ELISA units		
	treatment	prebleed	Week 3
546	water		22525
547	water		8939
548	water		11885
549	water		5121
550	water		37770
geometric mean			10521
pooled prebleed		11	
581	nair©		34222
582	nair©		45674
583	nair©		50224
584	nair©		27270
585	nair©		21832
geometric mean			38251
pooled prebleed		15	

#### Example 4.

Further studies were conducted to evaluate the effect of chemical penetration enhancement using a keratinolytic formulation (such as a salicylate). BALB/c mice 6 to 8 weeks of age were anesthetized and shaved as described in the “immunization procedure.” Twenty-four hours later, the backs of the “water” group were wiped with a gauze pad saturated in water and a pool of water was placed on the back.

Approximately 5 minutes later, any excess water was removed and 25 µg of CT (50 µl of a 0.5 mg/ml solution) was applied to the back. Alternatively, twenty-four hours after shaving, the backs of the “salicylate/water” group were treated with a gauze pad saturated with a 10% salicylate suspension (1 tablet (325 mg) Certified brand aspirin dissolved in 3.25 ml deionized water). Such treatment can continue for from about 0.1 to 30 minutes preferably about 20 minutes and more preferably about 10 minutes. Approximately 10 minutes later any remaining solution was blotted off, the backs of the animals were hydrated with water for 5 minutes, followed by removal of excess water, and then topical application of 25 µg of CT. Removal of excess antigen was conducted as described in the “immunization procedure.”

The anti-CT antibody titers were determined using ELISA as described above for “ELISA IgG (H+L)” 2 weeks after a single immunization. The results are shown in Table 4. While CT was clearly immunogenic in both groups, the geometric mean titer in the salicylate treated group was 4 fold higher and the titers were more consistent among the latter animals as compared with the “water” animals. Thus it appears that chemical disruption of the skin surface with salicylate enhances delivery of antigen by the transcutaneous route.

Table 4. Enhancement of transcutaneous immunization by chemical penetration enhancement: Anti-CT titers in mice that had the skin treated with salicylate (aspirin) before application of the antigen.

Animal #	anti-CT IgG (H+L) ELISA units		
	treatment	prebleed	week 2
741	water		272
742	water		not available
743	water		456
744	water		443
745	water		1395
geometric mean pooled prebleed		7	526
756	salicylate/water		2279
757	salicylate/water		4581
758	salicylate/water		4658
759	Salicylate/water		2771
760	Salicylate/water		593
geometric mean pooled prebleed		36	2402

### Example 5.

To assess the role of physical/mechanical penetration enhancement, an abrasive, in the form of a common emory board, was used to remove a portion of the stratum corneum. BALB/c mice 6 to 8 weeks of age were anesthetized and shaved as described in the “immunization procedure.” Twenty four hours later, the backs of the animals were either wiped with a gauze pad saturated in water “water” or brushed 10 times with a medium grain emory board “emory board.” and then wiped with a gauze pad saturated in water. Approximately five minutes after the water treatment, any excess water was removed and 20 µg of CT (100 µl of a 0.2 mg/ml) solution applied to the back. Removal of excess antigen was conducted as described in the “immunization procedure.”

The anti-CT antibody titers were determined using ELISA as described above for “ELISA IgG (H+L)” 3 weeks after a single immunization. The results are shown in Table 5. While CT was clearly immunogenic in both groups, the geometric mean titer in the emory board treated group was 10 fold higher and the titers were more consistent among the latter animals as compared with the “water” animals. Thus it appears that physical disruption of the outer surface of the skin with an emory board enhances delivery of antigen by the transcutaneous route. This can be differentiated from techniques that seek to pierce the skin and deliver antigen through the skin, such as in subcutaneous, intradermal or intramuscular injection.

This simple device could be replaced by other physical disrupting devices to deliver antigens and adjuvants into the epidermis such as microneedles that are of length to disrupt only the stratum corneum or superficial epidermis, devices used for TB tine testing, gas powered guns which do not penetrate the dermis, adhesive tape for tape stripping, or other barrier disruption devices known to disrupt only the stratum corneum or superficial epidermis.



Table 5. Enhancement of transcutaneous immunization by physical penetration enhancement: Anti-CT titers in mice that had the skin treated with an emory board before application of the antigen.

Animal #	anti-CT IgG (H+L) ELISA units		
	treatment	prebleed	week 3
7146	water		1275
7147	water		69
7148	water		7420
7149	water		6025
7150	water		388
geometric mean			1088
pooled prebleed		7	
7151	emory board		6632
7152	emory board		9380
7153	emory board		31482
7154	emory board		11142
7155	emory board		11761
geometric mean			12074
pooled prebleed		9	

### Example 6.

Another means of physical/mechanical penetration enhancement was employed using an abrasive pad to remove a portion of the stratum corneum and allow access to the underlying epidermis. BALB/c mice 6 to 8 weeks of age were anesthetized and shaved as described in the “immunization procedure”. Twenty four hours later, the backs of the animals were either wiped with a gauze pad saturated in water, “water”, or wiped with a gauze pad saturated in water followed by rubbing for 10 seconds with a nylon sponge (buf puf ©) to remove the outermost layers of the stratum corneum, “buf-puf ©”. Excess water was removed from the backs of the “water” group and then 20 µg of CT (100 µl of a 0.2 mg/ml solution) was applied to the backs of all animals. Removal of excess antigen was conducted as described in the “immunization procedure.”

The anti-CT antibody titers were determined using ELISA as described above for “ELISA IgG (H+L)” 3 weeks after a single immunization. The results are shown in Table 6. While CT was clearly immunogenic in both groups, the geometric mean titer in the buff puff treated group was 2 fold higher and the titers among individual animals were more consistent among the latter animals compared with the “water” animals. Thus it appears that physical disruption of the skin surface with a buff-puf © enhances delivery of antigen by the transcutaneous route.

This simple device could be replaced by other physical penetration devices to deliver antigens and adjuvants into the epidermis such as a needle and tuberculin syringe used for intradermal injection, microneedles that are of length to penetrate only the stratum corneum or superficial dermis, devices used for TB tine testing, abrading patches which have dissolvable crystals such as sucrose or sodium chloride or biodegradable polymers that are impregnated into the patch and rubbed on the skin before securing the patch with antigen either contained in the crystal or in the matrix, gas powered guns, adhesive tape for tape stripping, or other devices known to penetrate only into the epidermis or superficial dermis.

Table 6. Enhancement of transcutaneous immunization by physical penetration enhancement: Anti-CT titers in mice that had the skin treated with an abrasive pad (such as, for example, buf-puf ©) before application of the antigen.

Animal #	anti-CT IgG (H+L) ELISA units		
	treatment	prebleed	week 3
7146	water		1275
7147	water		69
7148	water		7420
7149	water		6025
7150	water		388
geometric mean pooled prebleed		7	1088
7166	buf puf©		5376
7167	buf puf ©		2319
7168	buf puf ©		1209
7169	buf puf ©		2871
7170	buf puf ©		2785
geometric mean pooled prebleed		8	2607

Example 7.

Transcutaneous immunization with bacterial ADP-ribosylating exotoxins such as CT and LT appear to provide significant 'danger' signals to the immune system stimulating a potent immune response. Such compounds act as adjuvants. It was a surprise to find that simple mixtures of such adjuvants placed on the skin in a manner that hydrates the skin, resulting in potent immune responses. This was described in earlier patents (PCT/US97/21324). However, given that an adjuvant such as CT (86KD) can act as an adjuvant on the skin, it would be expected that other adjuvants, particularly those based on bacterial products or motifs, would be stimulatory when placed on the skin in a manner that hydrates the skin and/or with the use of penetration enhancers.

We used bacterial DNA to confirm that this expectation is correct. BALB/c mice 6 to 8 weeks of age were shaved and anesthetized as described above for the "immunization procedure". On the day of immunization the backs of the mice were wiped with isopropanol to enhance penetration. After the alcohol had evaporated (approximately 5 minutes), 100 µl of phosphate buffered saline (PBS) containing 100 µg of DNA (CpG1 or CpG2) and 100 µg of diphtheria toxoid (DT) was applied to the back for 90 to 120 minutes. Oligonucleotides were synthesized by Oligos Etc with a phosphorothioate backbone to improve stability. Removal of excess antigen was conducted as described in the "immunization procedure." The immunization was repeated 4 and 8 weeks later. Ten weeks after the primary immunization the animals were bled and the anti-DT titers determined using an ELISA as described above for "ELISA IgG (H+L)". The results are shown in Table 7A.

Sub<sub>C1</sub> Co-administration of DT and a control DNA sequence (CpG2: TCCAATGAGCTTCCTGAGTCT) failed to induce a detectable rise in the anti-DT titers. In contrast, addition of a DNA sequence containing an unmethylated CpG dinucleotide flanked by two 5' purines and two 3' pyrimidines (CpG1 (immunostimulatory DNA): TCCATGACGTTTCCTGACGTT) resulted in a detectable increase in the serum anti-DT IgG titer in 5 of 5 animals. Thus it appears that bacterial DNA containing appropriate motifs such as CPGs (6KD) can be used as adjuvant to enhance delivery of antigen through the skin for induction of antigen specific antibody responses.

Table 7A. Adjuvant activity of bacterial DNA applied to the skin using penetration enhancement: humoral immune response.

Anti-DT IgG (H+L) ELISA units		
Animal #	adjuvant/antigen prebleed	week 10
7261	CpG1/DT	1171
7262	CpG1/DT	22750
7263	CpG1/DT	4124
7264	CpG1/DT	126
7265	CpG1/DT	115
Geometric mean pooled prebleed	6	1096
7266	CpG2/DT	19
7267	CpG2/DT	12
7268	CpG2/DT	5
7269	CpG2/DT	5
7270	CpG2/DT	11
geometric mean pooled prebleed	5	9

Sub<sup>5</sup>  
C<sub>2</sub>

The transcutaneous effect of transcutaneous immunization can also be detected by T-cell proliferation. BALB/c mice 6 to 8 weeks of age were shaved and anesthetized as described above for the "immunization procedure". On the day of immunization the backs of the mice were wiped with isopropanol. After the alcohol had evaporated (approximately 5 minutes), 100 µl of phosphate buffered saline (PBS) containing 100 µg of DNA (CpG1 or CpG2) and 100 µg of diphtheria toxoid (DT) was applied to the back for 90 to 120 minutes. Oligonucleotides were synthesized by Oligos Etc with a phosphorothioate backbone to improve stability. Removal of excess antigen was conducted as described in the "immunization procedure." The immunization was repeated 4 and 8 weeks later. Twelve weeks after the primary immunization draining (inguinal) LNs were removed and pooled from five immunized animals. The capacity

to proliferate in response to media or antigen (DT) was assessed in a standard 4 day proliferation assay using 3-H incorporation as a readout. The results are shown in Table 7B. Co-administration of DT and a DNA sequence containing an unmethylated CpG dinucleotide flanked by two 5' purines and two 3' pyrimidines (CpG1  
5 (immunostimulatory DNA): *TCCATGACGTTTCCTGACGTT*) resulted in a detectable increase in the antigen specific proliferative response. Thus it appears that bacterial DNA containing appropriate motifs can be used as adjuvant to enhance delivery of antigen through the skin for induction of proliferative responses.

Table 7B. Adjuvant effect of bacterial DNA applied to the skin: LN cell proliferation

antigens applied in vivo	proliferation (cpm) 3-H incorporation in vitro to antigens	
	media	DT
normal LN	339	544
CpG1/DT	1865	5741

#### Example 8.

Transcutaneous immunization with bacterial ADP-ribosylating exotoxins such as CT and LT appear to provide significant 'danger' signals to the immune system stimulating a potent immune response. Such compounds act as adjuvants. It was a surprise to find that simple mixtures of such adjuvants placed on the skin in a manner that hydrates the skin, resulting in potent immune responses. This was described in earlier patents (PCT/US97/21324). However, given that an adjuvant such as CT can act as an adjuvant on the skin, it would be expected that other adjuvants, would be stimulatory when placed on the skin in a manner that hydrates the skin. Genetically altered toxins were used to confirm this expectation. BALB/c mice 6 to 8 weeks of age were anesthetized, shaved, and immunized as described in the "immunization procedure". The animals were boosted 3 and 5 weeks after the primary immunization and serum collected 2 weeks after the final immunization. The adjuvants used were the genetically altered toxins; LTK63, an enzymatically inactive LT derivative, and LTR72, an LT derivative which retains 0.6% of the enzymatic activity. Diphtheria toxoid (DT) 100 µg was used as antigen.

Anti-DT antibody titers were determined using ELISA as described above for "ELISA IgG (H+L)". The results are shown in Table 8. Anti-DT titers were clearly elevated in serum from animals immunized with either LTR63 or LTR72 and DT when compared with titers in serum collected prior to immunization (prebleed). Thus it appears that genetically detoxified mutants of heat labile enterotoxin (LT) can be used as adjuvants on the skin.

Table 8. Use of genetically altered toxins, LTK63 and LTR72, as adjuvants on the skin.

Animal #	anti-DT IgG (H+L) ELISA units		
	adjuvant / antigen	prebleed	week 7
653	LTK63 / DT		20228
654	LTK63 / DT		not available
655	LTK63 / DT		342
656	LTK63 / DT		2445
657	LTK63 / DT		< 100
geometric mean pooled prebleed		<100	1140
663	LTR72 / DT		12185
664	LTR72 / DT		10917
665	LTR72 / DT		151
666	LTR72 / DT		2057
667	LTR72 / DT		50923
geometric mean pooled prebleed		<100	4620



### Example 9.

Another class of compounds, cytobines which are known to act as adjuvants illustrate the principle that adjuvants in general could be expected to act in a fashion similar to cholera toxin.  $\text{TNF}\alpha$  is also known to be a Langerhan cell activating compound.

BALB/c mice 6 to 8 weeks of age were shaved and anesthetized as described above for the "immunization procedure". On the day of immunization the backs of the mice were wiped with isopropanol. After the alcohol had evaporated (approximately 5 minutes), 100  $\mu\text{l}$  of phosphate buffered saline (PBS) containing 0.83  $\mu\text{g}$  TNF-alpha (recombinant mouse TNF-alpha, Endogen), IL-2 (1  $\mu\text{g}$  recombinant mouse IL-2 (Sigma)) or mock adjuvant (CpG2) was applied to the skin on the back with 100  $\mu\text{g}$  of diphtheria toxoid (DT) for 90 to 120 minutes. Oligonucleotides were synthesized by Oligos Etc with a phosphorothioate backbone to improve stability Removal of excess antigen was conducted as described in the "immunization procedure." The immunization was repeated 4 and 8 weeks later. Ten weeks after the primary immunization the animals were bled and the anti-DT titers determined using an ELISA as described above for "ELISA IgG (H+L)". The results are shown in Table 9.

Co-administration of DT and a mock adjuvant (CpG2) failed to induce a detectable rise in the anti-DT titers. In contrast, topical application of TNF-alpha (0.8  $\mu\text{g}$ ) resulted in a detectable increase in the serum anti-DT IgG titer in 3 of 5 animals when compared with either anti-DT titers in the mock adjuvant treated mice or sera collected prior to immunization (prebleed). Similarly, topical application of IL-2 (1  $\mu\text{g}$ ) resulted in a detectable increase in the serum anti-DT IgG titer in 4 of 5 animals when compared with either anti-DT titers in the mock adjuvant treated mice or sera collected prior to immunization (prebleed). Thus it appears that the cytokines such as IL-2 and TNF-alpha can be used as an adjuvant on the skin and that langerhans cell activating compounds can be used for transcutaneous immunization.

Table 9. Adjuvant activity of the cytokine TNF-alpha applied to the skin.

Animal #	adjuvant/antigen	Anti-DT IgG (H+L) ELISA units	
		prebleed	week 10
7326	TNF-alpha/DT		1808
7327	TNF-alpha/DT		830
7328	TNF-alpha/DT		7
7329	TNF-alpha/DT		1477
7330	TNF-alpha/DT		7
geometric mean pooled prebleed	1		159
7331	IL-2/DT		13
7332	IL-2/DT		111
7333	IL-2/DT		345
7334	IL-2/DT		49
7335	IL-2/DT		35
geometric mean pooled prebleed	2		61
7266	CpG2/DT		19
7267	CpG2/DT		12
7268	CpG2/DT		5
7269	CpG2/DT		5
7270	CpG2/DT		11
geometric mean pooled prebleed	5		9

## Example 10.

The B-subunit of cholera toxin is another class of adjuvants lack the A-subunit and therefore ADP-ribosyltransferase activity of CT. As such CTB represents an  
5 adjuvant that is unique and may be useful as it is not toxic when ingested.

C57BL/6 mice 6 to 8 weeks of age were anesthetized and shaved as described in the "immunization procedure". On the day of immunization the backs of the mice were wiped with isopropanol. After the alcohol had evaporated (approximately 5 minutes), 100 µl of phosphate buffered saline (PBS) containing 100 µg of purified  
10 cholera toxin B subunit (CTB) and/or 100 µg of diphtheria toxoid (DT) was applied to the back for 90 to 120 minutes. Removal of excess antigen was conducted as described in the "immunization procedure." The immunization was repeated 4 and 8 weeks later. Ten weeks after the primary immunization the animals were bled and the anti-DT titers determined using an ELISA as described above for "ELISA IgG  
15 (H+L)". The results are shown in Table 10.

Anti-DT titers were clearly elevated in serum from animals immunized with CTB and DT when compared with titers in serum from animals treated with DT alone or those in prebleed serum samples as shown in Table 10. Thus it appears that purified CTB can be used as an adjuvant on the skin.

Table 10. Use of purified cholera toxin B subunit from *V. cholerae* as an adjuvant on the skin.

Animal #	Anti-DT IgG (H+L) ELISA units		
	adjuvant / antigen	prebleed	week 10
51	DT		11
52	DT		7
53	DT		4
54	DT		8
55	DT		7
geometric mean			7
pooled prebleed		4	
81	CTB/DT		14880
82	CTB/DT		371
83	CTB/DT		14810
84	CTB/DT		108
85	CTB/DT		27
geometric mean			751
pooled prebleed		5	

Example 11.

Adjuvants that are structurally different are likely to exert their enhancement by different effects. Adjuvants that induce their effects by different mechanisms may have either additive or synergistic effects on enhancing the immune response. We found that the use of two adjuvants simultaneously augmented the response to transcutaneous immunization compared to the individual adjuvants alone.

BALB/c mice 6 to 8 weeks of age were shaved and anesthetized as described above for the "immunization procedure". On the day of immunization the backs of the mice were wiped with isopropanol. After the alcohol had evaporated (approximately 5 minutes), 100 µl of phosphate buffered saline (PBS) containing 100 µg of immunostimulatory DNA (CpG1) and/or cholera toxin (CT) 100 µg was applied to the back with 100 µg of a soluble leishmanial antigen extract (SLA) for 90 to 120 minutes. SLA is an antigen extract prepared at Walter Reed Army Institute of Research by centrifugal isolation of the soluble proteins in a sonicate of *Leishmania major* promastigotes) extract for 90 to 120 minutes. Removal of excess antigen was conducted as described in the "immunization procedure." The immunization was repeated 4 and 8 weeks later. Twelve weeks after the primary immunization draining (inguinal) LNs were removed and pooled from two immunized animals. The capacity to proliferate in response to media or antigen (SLA) was assessed in a standard 4 day proliferation assay using 3-H incorporation as a readout. The results are shown in Table 11.

Sub  
13 Co-administration of SLA and CpG1 (immunostimulatory DNA containing an unmethylated CpG dinucleotide flanked by two 5' purines and two 3' pyrimidines - TCCATGACCGTTCTGACGTT) or CT resulted in a detectable increase in the antigen specific proliferative response. However, the antigen (SLA) specific proliferative response was approximately 20 times higher in lymph node cell cultures from animals exposed simultaneously to both CpG1 and CT as compared to cultures derived from animals exposed to either adjuvant alone. Thus it appears that bacterial DNA containing appropriate motifs synergizes with ADP ribosylating exotoxins such as CT as adjuvants on the skin to induce higher immune responses than to either adjuvant alone.

Table 11. Synergy between immunostimulatory DNA and ADP ribosylating exotoxin (CT) as adjuvants when applied to the skin

<u>proliferation (cpm) 3-H incorporation in vitro to antigens</u>		
<u>substances applied in vivo</u>	<u>media</u>	<u>SLA</u>
normal LN	180	219
SLA	200	159
SLA/CpG1	1030	2804
SLA/CT	232	2542
SLA/CpG1/CT	2232	47122

#### Example 12.

Transcutaneous immunization induces potent immune responses when used as a method of delivery alone. We also have found that transcutaneous immunization can be used in sequence with other routes of delivery to stimulate an immune response.

BALB/c mice 6 to 8 weeks of age On day 0 both groups of animals received a 50 µl intramuscular (im.) injection of DT (5 µg) mixed with alum (Rehydrogel - 25 µg in NaCl) into the hind thigh. Eight and 16 weeks later mice in the im/tc/tc group were shaved, anesthetized and immunized by the transcutaneous route as described above for the "immunization procedure" using 100 µg cholera toxin as adjuvant and 100 µg diphtheria toxoid as antigen. The immunization solution was applied to the back for 90 to 120 minutes. Removal of excess antigen was conducted as described in the "immunization procedure." Twenty two weeks after the primary immunization the animals were bled and the anti-DT titers determined using an ELISA as described above for "ELISA IgG (H+L)". The results are shown in Table 12.

A single im. injection of 5 µg of DT induced a detectable rise in the serum anti-DT titers as compared with titers in sera collected from the same animals prior to immunization (prebleed). Boosting of the im. primed animals using the transcutaneous immunization method resulted in an 60 fold rise in the geometric mean titer and clearly all transcutaneously boosted animals had higher anti-DT titers than those observed in the im. primed group. Thus transcutaneous immunization can be

used to boost antigen specific titers in mice in which the primary immunization with the antigen was by the i.m. route. We have also found that im primed animals can be boosted by transcutaneous immunization (TCI). Various combinations of TCI priming or boosting with other routes and schedules can be visualized including oral,

5 buccal, nasal, rectal, vaginal, intradermal, by gun or other means of delivery. Additionally, antigens may differ in route and composition including protein alternating with glycoprotein, subunit with holotoxin, DNA priming followed by protein, nucleic acid by im followed by nucleic acid by TCI. Transcutaneous immunization may be used to boost children primed in infancy or adults primed in

10 childhood. The ease of delivery may enhance the efficacy vaccines such as the influenza vaccines by allowing multiple boosts using a patch.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212

Table 12. Boosting of im. primed animals using the transcutaneous immunization method

Anti-DT IgG (H+L) ELISA units				
Animal #	adjuvant/antigen	Route of administration	prebleed	week 22
8563	DT	im.		54227
8564	DT	im.		11833
8565	DT	im.		106970
8566	DT	im.		10830
8567	DT	im.		4003
geometric mean pooled prebleed			20	19711
8568	DT/ct+dt/ct+dt	im./tc./tc.		628838
8569	DT/ct+dt/ct+dt	im./tc./tc.		2035507
8570	DT/ct+dt/ct+dt	im./tc./tc.		1164425
8571	DT/ct+dt/ct+dt	im./tc./tc.		not available
8572	DT/ct+dt/ct+dt	im./tc./tc.		1263138
geometric mean pooled prebleed			10	1171368
8558	DT /DT/DT	im./im./im.		not available
8559	DT /DT/DT	im./im./im.		542669
8560	DT /DT/DT	im./im./im.		770150
8561	DT /DT/DT	im./im./im.		545894
8562	DT /DT/DT	im./im./im.		671898
geometric mean pooled prebleed			15	625721



### Example 13.

Because TCI appears to stimulate the immune system in such a potent fashion, it is possible that an adjuvant place on the skin at one site may act as an adjuvant for an antigen placed at another site. BALB/c mice 6 to 8 weeks of age were anesthetized and shaved as described in the “immunization procedure”. Animals were not ear tagged but kept in cages labeled A, C or G. On the day of immunization the dorsal surface of the mouse ear was treated by gently rubbing the outer skin surface with a cotton-tipped applicator containing 70% isopropanol. After 5 minutes the excess water was blotted from water-treated ears and adjuvant (CT 50 µg) and/or antigen (100 µg of bovine serum albumin (BSA) was applied to the left or right ear surface (described in table) in 50 µl phosphate buffered saline. At two and a half hours, the ears were rinsed and blotted dry twice. Mice were boosted in a similar fashion four and eight weeks later. Twelve weeks after the primary immunization the animals were bled and the anti-BSA titers determined using an ELISA as described above for “ELISA IgG (H+L)”. The results are shown in Table 13.

Application of BSA alone to the skin was poorly immunogenic with only 1 of 5 animals developing an ELISA titer above 100 EU. In contrast, 9 of 9 animals receiving CT and BSA on the skin developed titers above 100 EU. Of the animals receiving antigen and adjuvant, the mice given the materials at the same site (left ear) developed higher (10 fold) anti-BSA titers than animals receiving antigen and adjuvant in separate (left and right) ears. However, animals receiving antigen on one ear and adjuvant on another ear developed an anti-BSA immune response that was approximately 30 times higher than animals given BSA alone. Thus, antigen and adjuvant may be topically applied during TCI at different sites to elicit a humoral immune response. This immunostimulation may be expected to occur with antigen delivered by other routes and schedules can be visualized including oral, buccal, nasal, rectal, vaginal, intradermal, by gun or other means of delivery. Additionally, adjuvants may be used with nucleic acid immunization to enhance the response. Such a delivery may not need to be simultaneous to enhance the immune response. For example, an im injection of plasmid DNA may be followed later by transcutaneous administration of adjuvant. Immunostimulation by CT, LT, TNF $\alpha$ , CpGs or similar

adjuvants is a surprising result because it has been thought that molecules 500 daltons could not pass through the skin.

5 Table 13. Delivery of antigen and adjuvant at the same or distal sites on the skin with penetration enhancement.

Anti-BSA IgG (H+L) ELISA units			
Animal #	adjuvant / antigen	prebleed	week 12
group G	BSA left ear		240
group G	BSA left ear		99
group G	BSA left ear		40
group G	BSA left ear		not available
group G	BSA left ear		15
Geometric mean pooled prebleed		6	61
group C	CT/BSA left ear		16418
group C	CT/BSA left ear		24357
group C	CT/BSA left ear		13949
group C	CT/BSA left ear		70622
group C	CT/BSA left ear		not available
Geometric mean pooled prebleed		3	25053
group A	CT left/BSA right ear		106
group A	CT left/BSA right ear		23806
group A	CT left/BSA right ear		1038
group A	CT left/BSA right ear		1163
group A	CT left/BSA right ear		8696
Geometric mean pooled prebleed		15	1939

#### Example 14. Transcutaneous Immunization in Humans

The invention can be practiced using a suitable vehicle or carrier. For example a patch can be used as such a vehicle and can be treated with the formulation of the invention or can be used to cover the area of the skin which has been treated with the formulation of the invention. A suitable patch can be fabricated from, for example, cotton, nylon, rayon, polyester or combinations thereof. Such patches can be provided with an adhesive or non-adhesive backing. Patches with a non-adhesive backing can be secured to the animal by non-adhesive means, such as, for example, wrapping.

Suitable backing can be fabricated from materials such as, for example, silicon, acrylate or rubber. Other vehicles or carriers which can be used include those listed above; such as for example, powders, oils, water, cream and the like. To evaluate the potential for TCI in humans, a Phase I trial was conducted using LT to attempt to induce serum anti-LT antibodies. Six volunteers received a dose of 500 ug of LT, a dose similar to oral adjuvant doses used for a cholera vaccine (1 mg CTB). LT was produced under GMP conditions at the Swiss Serum and Vaccine Institute (Berne, Switzerland) and was provided by Oravax Inc., Cambridge, MA. The volunteers received 500 ug of LT mixed in 500 ul of sterile saline which was absorbed on a 2in<sup>2</sup> cotton gauze pad with polyvinyl backing and covered by a 4x4 in<sup>2</sup> Tegaderm<sup>TM</sup> dressing. The immunization was conducted by placing the patch on unmanipulated skin for 6 hours after which the site was thoroughly rinsed with 500 ml of sterile saline. Individuals were similarly reimmunized after 12 weeks. Volunteers were examined on days 1, 2, 3 and 7 for signs of inflammation at the site of immunization and interviewed for symptoms related to the vaccination. The immunization was conducted by placing the patch on unmanipulated skin for 6 hours after which the patch was removed and the site was thoroughly rinsed with saline. Individuals were reimmunized after 12 weeks. No adverse reactions were seen, either systemically or at the site of immunization after the first or second immunization. Anti-LT IgG titers were determined as previously described. Results are reported in ELISA units (EU) which are defined as the inverse dilution of sample that yields an OD of 1.0. Anti-LT IgA was determined in the same manner as anti-LT IgG using goat anti-human IgA( $\alpha$ )-HRP (Kirkegaard and Perry, Gaithersburg, MD) against a standard IgA curve made using human IgA (ICN). As shown in Table 14, 6 of 6 individuals responded

by producing a rise in serum anti-LT IgG or IgA antibodies, defined as a four-fold increase in antibody titers. The mean fold rise in anti-LT IgG was 10.2 and the mean fold rise in serum anti-LT IgA was 7.2. Biopsies of the immunization site and contralateral arm showed no signs of inflammation of the skin. These results confirm that TCI can be conducted in humans without skin irritation or inflammation.

Suitable patch materials have been previously described. In general the patch may consist of, for example, a pressure sensitive dressing, a matrix for or absorbant layer to carry the vaccines and adjuvant, a vaccine impermeable backing and a release liner. This and other suitable patch examples are described in U.S. Patent Nos. 4,915,950 and 3,734,097.

Patches can be fabricated to include woven and non-woven matrices of materials to include, for example, polyester/cellulose, polypropylene, polyester, polyester/rayon and the like.

Examples of non-woven patch matrices can include:

#### BBA Nonwovens

- a. Grade# 1313290, wet laid non-woven, composition = polyester/cellulose, weight (gsy) = 35.4, weight (gsm) = 42.3, thickness (mils) = 7.9, tensile MD = 3.4, tensile CD = 2.4.
- b. Grade# 2006086, thermal bond non-woven, composition = polypropylene, weight (gsy) = 16.0, weight (gsm) = 19.1, thickness (mils) = 10.2, tensile MD = 3.3, tensile CD = 0.7.
- c. Grade# 149146, thermal bond non-woven, composition = polyester, weight (gsy) = 25.6, weight (gsm) = 30.6, thickness (mils) = 6.5, tensile MD = 5.3, tensile CD = 0.9.
- d. Grade# 149020, thermal bond non-woven, composition = polyester/rayon, weight (gsy) = 30.5, weight (gsm) = 36.4, thickness (mils) = 13.2, tensile MD = 5.5, tensile CD = 1.1.
- e. Grade# 140-027, hydroentangled nonwoven, composition = polyester/rayon, weight (gsy) = 28.0, weight (gsm) = 33.5, thickness (mils) = 22.4,

tensile MD = 10.4, tensile CD = 3.8.

Pressure sensitive adhesive that can be used in the present invention include, for example adhesive based on acrylate, silicone, rubber and the like.

5

**Table 14** Mean fold rise in human anti-LT IgG and IgA

Volunteer #	4 week IgG	12 week IgG	16 week IgG
13	15.2	9.5	12.5
14	1.4	1.6	1.7
15	11.7	15.0	12.9
16	1.3	0.7	16.0
17	12.5	51.9	58.6
18	1.3	2.1	4.3
Mean rise IgG	4.2	5.0	10.2
Volunteer #	4 week IgA	12 week IgA	16 week IgA
13	7.2	4.1	10.1
14	4.9	4.3	4.3
15	4.9	5.7	4.5
16	1.4	1.3	7.0
17	15.3	29.4	28.1
18	1.3	1.5	3.5
Mean rise IgA	4.1	4.2	7.2

Example 15.

10 LT has been shown to be effective for immunizing humans by the transcutaneous route. We also have found that LT acts as an adjuvant for TCI. C57BL/6 mice 6 to 8 weeks of age were anesthetized and shaved as described in the "immunization procedure". On the day of immunization the backs of the mice were wiped with isopropanol. After the alcohol had evaporated (approximately 5 minutes),  
15 100 µl of phosphate buffered saline (PBS) containing 100 µg of heat labile enterotoxin (LT) and/or 100 µg of diphtheria toxoid (DT) was applied to the back for 90 to 120 minutes. Removal of excess antigen was conducted as described in the "immunization procedure." The immunization was repeated 4 and 8 weeks later. Ten weeks after the primary immunization the animals were bled and the anti-DT titers determined using

an ELISA as described above for "ELISA IgG (H+L)". The results are shown in Table 15.

Anti-DT titers were clearly elevated in serum from animals immunized with LT and DT when compared with titers in serum from animals treated with DT alone or those in prebleed serum samples. Thus it appears that heat labile enterotoxin (LT) can be used as an adjuvant on the skin.

Table 15. Use of heat labile enterotoxin (LT) from *E. coli* as an adjuvant on the skin.

Anti-DT IgG (H+L) ELISA units			
Animal #	adjuvant / antigen	prebleed	week 10
51	DT		11
52	DT		7
53	DT		4
54	DT		8
55	DT		7
geometric mean			7
pooled prebleed		4	
71	LT/DT		7126
72	LT/DT		46909
73	LT/DT		669
74	LT/DT		8480
75	LT/DT		1598
geometric mean			4970
pooled prebleed		5	

Example 16.

To assess the role of physical/mechanical penetration enhancement, the superficial layers of the stratum corneum were removed by tape stripping. Tape stripping is an intervention well known in the art to remove the outer layer of the stratum corneum. C57BL/6 mice 6 to 8 weeks of age were anesthetized and shaved as described in the "immunization procedure." Twenty four hours later, CT (25 µg) was applied to the backs of the mice in 50 µl of phosphate buffered saline; "none" intervention group. Alternatively, the skin on the backs of a second group of animals was subjected to mild tape stripping; "tape stripping" intervention group. The tape stripping procedure was accomplished by applying cellophane scotch-tape to the backs, allowing bonding to the skin surface for 3 minutes, followed by gentle removing of the tape. The bonding/removal steps were repeated 3 times. CT (25 µg) was then applied to the backs of the mice in 50 µl of phosphate buffered saline. Antigen remained on the backs for approximately 1.5 hrs at which time removal of excess antigen was conducted as described in the "immunization procedure."

The anti-CT antibody titers were determined using ELISA as described above for "ELISA IgG (H+L)" using sera collected 11 days after the primary immunization. The results are shown in Table 16. CT was immunogenic in both groups as compared to sera collected from the same animals prior to immunization (prebleed). However, the geometric mean titer in the tape stripped group was 100 fold higher and the titers were more consistent among the latter animals as compared with the "none" animals. Thus it appears that physical disruption of the skin surface using tape stripping enhances delivery of antigen by the transcutaneous route.

This simple device could be replaced by other physical penetration devices to deliver antigens and adjuvants into the epidermis such as a needle and tuberculin syringe used for intradermal injection, microneedles that are of length to penetrate only the stratum corneum or superficial dermis, devices used for TB tine testing, gas powered guns, adhesive tape for tape stripping, or other devices known to penetrate only into the epidermis or superficial dermis. Tape stripping devices could be used in conjunction with other penetration enhancers. Tape stripping devices may be used in conjunction with a marker to delineate the site for patch placement, and may be dispersed in a roll or in individual units.



Table 16. Enhancement of transcutaneous immunization by physical penetration enhancement: Anti-CT titers in mice that had the skin stripped using cellophane tape before application of the antigen.

Animal #	intervention	anti-CT IgG (H+L) Elisa units	
		prebleed	dav 11
976	none		155
977	none		4
978	none		4
979	none		31
980	none		23
geomean prebleed		2	16
986	tape strip		10702
987	tape strip		1285
988	tape strip		5832
989	tape strip		997
990	tape strip		782
geomean prebleed		3	2990

#### Example 17.

Nucleic acids such as plasmid DNA or RNA can be used to induce and immune response and are well known in the art. The use of Nucleic acids in transcutaneous immunization was described in previous patents (PCT/US97/21324). The use of nucleic acids (also known as genetic immunization) on the skin with penetration enhancement techniques is illustrated in the following example.

C57BL/6 mice 6 to 8 weeks of age were anesthetized and shaved as described in the "immunization procedure". For the "NP DNA" group the mice were wiped with isopropanol, after the alcohol had evaporated (approximately 10 minutes), the backs were hydrated with water using a saturated gauze pad. Approximately 10 minutes later the any excess water was blotted off and a 100 µg of a DNA plasmid (pCMV-NP) encoding for influenza nucleoprotein was applied to the back in 100 µl of

saline. A second group of NP-DNA mice were subjected to the same immunization protocol except that their backs were tape stripped 3x prior to alcohol swabbing; "NP DNA – tape stripping". The tape stripping procedure was accomplished by applying cellophane scotch-tape to the backs, allowing bonding to the skin surface for 5

5 minutes, followed by gentle removing of the tape. A third group of mice was engaged in the tape stripping / immunization protocol described and 100 µg of the adjuvant heat labile enterotoxin (LT) was included in the immunization solution. Sixteen days after the primary immunization the animals were bled and the anti-influenza NP titers determined using an ELISA as described above for "ELISA IgG (H+L)". The results are shown in Table 17.

10 Anti-influenza titers were determined using a split virus antigen (Fluzone) preparation to coat the ELISA plates. ELISA titers were determined in 5 individual animals and the mean optical density reading for each group is shown. All three immunization groups developed anti-influenza titers as compared with titers in serum collected from the same animals prior to immunization (prebleed). As compared with the NP DNA alone group, tape stripping prior to immunization enhanced the anti-Influenza titer in all three serum dilutions tested (1:100, 1:200, 1:400) and addition of an adjuvant (LT) further enhanced this response. Thus, DNA can be used on the skin to induce immune responses to vaccine antigens and its effectiveness can be enhanced by the addition of adjuvants and penetration enhancement such as tape stripping.

Table 17. Immuogenicity of DNA applied as antigen on the skin using alcohol penetration enhancement.

Antigen/adjuvant	intervention	Anti-INF IgG optical density (405 nm)			
		prebleed	day 16 post immunization		
		1:100	1:100	1:200	1:400
NP DNA	none	0.21	0.47	0.20	0.07
NP DNA	tape stripping	0.39	0.64	0.28	0.13
NP DNA / LT	tape stripping	0.39	0.87	0.38	0.13

### Example 18.

Transcutaneous immunization (TCI), because of the ease of delivery, allows the application to be given over different draining lymph nodes. This may have an additional advantage in that it enhances the immune response. Rabbits were anesthetized, shaved, and immunized as described above. Animals were immunized with 100 µg cholera toxin (CT) and 100 µg of influenza hemagglutinin (HA) at one site or two sites on the back. HA and CT were applied at 0, 3 and 5 weeks. Seven weeks after the primary immunization, the animals were bled and the anti-HA titers determined using an ELISA as described above for "ELISA IgG (H+L)". The results are shown in Table 18.

Anti-HA titers were elevated in serum from 10 of 10 animals immunized with CT and HA when compared with titers in serum from the same animals prior to immunization (prebleed). The geometric mean titer in the two site group was 3 fold higher than that in the one site group suggesting that antigen delivery at multiple sites may be used to enhance TCI. Thus, antigens can be delivered by TCI either at a single or multiple sites on the skin.

Table 18. Transcutaneous delivery of antigen in a single or multiple sites.

Animal	antigen/adjuvant	anti-HA IgG (ELISA units)		
		prebleed	7 weeks	geomean
1	CT/HA one site	< 25	1142	2596
2	CT/HA one site	< 25	9617	
3	CT/HA one site	< 25	2523	
4	CT/HA one site	< 25	2275	
5	CT/HA one site	< 25	1869	
6	CT/HA two sites	< 25	10348	8403
7	CT/HA two sites	< 25	18453	
8	CT/HA two sites	< 25	9778	
9	CT/HA two sites	< 25	15985	
10	CT/HA two sites	< 25	1404	

# Example 19

The variety of antigens which can be delivered by TCI is further illustrated by the use of a polysaccharide conjugate vaccine to induce anti-polysaccharide antibodies. BALB/c mice 6 to 8 weeks of age were anesthetized, shaved, and immunized as described in the “immunization procedure”. Mice were immunized with cholera toxin (CT) and *Haemophilus influenzae B* polysaccharide (Hib-PS) at 0, 3 and 5 weeks. Seven weeks after the primary immunization, the animals were bled and the anti-Hib-PS titers determined using an ELISA as described above for “ELISA IgG (H+L)”. The results are shown in Table 19.

Anti-Hib-PS titers were elevated in serum from 4 of 10 animals immunized with CT and Hib-PS when compared with titers in serum from the same animals prior to immunization (prebleed). Thus TCI can be used to induce anti-polysaccharide antigens through the skin. This is a common human use vaccine antigen and represents an important strategy for immunization.

Table 19. Delivery of a conjugated polysaccharide by transcutaneous immunization

ear tag #	antigen/adjuvant	anti-Hib PS IgG (µg/ml)	
		prebleed	7 weeks
1	CT/Hib-PS (100µg/100µg)	< 0.20	< 0.20
2	CT/Hib-PS (100µg/100µg)	< 0.20	< 0.20
3	CT/Hib-PS (100µg/100µg)	< 0.20	<u>1.68</u>
4	CT/Hib-PS (100µg/100µg)	< 0.20	< 0.20
5	CT/Hib-PS (100µg/100µg)	< 0.20	<u>1.86</u>
6	CT/Hib-PS (100µg/25µg)	< 0.20	<u>1.04</u>
7	CT/Hib-PS (100µg/25µg)	< 0.20	< 0.20
8	CT/Hib-PS (100µg/25µg)	< 0.20	< 0.20
9	CT/Hib-PS (100µg/25µg)	< 0.20	<u>6.30</u>
10	CT/Hib-PS (100µg/25µg)	< 0.20	< 0.20

## Example 20

### Transcutaneous immunization of mice with human use vaccine antigens

CT has been shown to act as adjuvant for transcutaneous immunization with single toxoids and BSA. We transcutaneously immunized mice with a variety of human-use

5 vaccine antigens, including a multivalent toxoid vaccine (tetanus and diphtheria toxoids), a yeast expressed recombinant protein(HIV p55 gag) and whole killed rabies viruses using CT as an adjuvant as shown in Table 20. BALB/c mice (n=5) were immunized and boosted twice as previously described (Glenn, G.M., Scharton-Kersten, T., Vassell, R., Matyas, G. & Alving, C.R. Transcutaneous immunization using bacterial ADP-ribosylating exotoxins as antigens and adjuvants. *Infect. Immun.* (in the press). Immunizing doses included 100/50/50 ug CT/TT/DT via TCI versus 3/1/1 for alum/TT/DT IM; 100/100 ug for LT+DT versus 100 ug of DT alone.

10 100/100 ug for CT /p55 via TCI versus 100 ug p55 alone. Mice immunized with 17 IE of killed rabies virus (n=10) had been primed intramuscularly 2x and then boosted transcutaneously (17 IE) after light alcohol swabbing of the skin and compared to 3x IM rabies immunization. Antibody levels against DT, TT, p55 and rabies were determined using ELISA as previously described (Grassi, M., Wandler, A. & Peterhans, E. Enzyme-linked immunoabsorbant assay for determination of antibodies to the envelope glycoprotein of rabies virus. *J.Clin.Microbiol.* 27, 899-902 (1989).

15 20 Miyamura, K., Tajiri, E., Ito, A., Murata, R. & Kono, R. Micro cell culture method for determination of diphtheria toxin and antitoxin titres using VERO cells. II.

Comparison with the rabbit skin method and practical application for seroepidemiological studies. *J. Biol. Stand.* 2, 203-209 (1974)). TCI resulted in similar increases in the antibody responses to TT and DT and the anti-DT

25 neutralization titers were comparable to that elicited by intramuscular immunization (IM). These data show that TCI may be used to <sup>induce</sup> ~~induce~~ immune response of comparable magnitude as those induced by existing immunization practices. TCI boosting of IM primed animals also resulted in a significant rise in anti-rabies titers in all 10 animals tested (0.53 to 1.03 IU, p<0.02, Student's t-test). Antibodies to the  
30 antigens DT and p55 administered without adjuvants were very low or undetectable, consistent with our previous observations that antigens are only weakly immunogenic when applied without adjuvant. LT also acted as adjuvant (Table 20) in a fashion

5 vaccine antigens.



Table 20.

Murine antibody responses to human-use vaccine antigens administered by TCI

Immunizing Antigen(s)for TCI	Antibody specificity	TCI (ELISA Units)	IM/alum (ELISA Units)
CT+TT+DT	Anti-DT	135,792 (86,552-146-759)	85,493 (24,675-238,904)
CT+TT+DT	Anti-TT	30,051 (13,863-53,174)	94,544 (74,928-113,408)
CT+TT+DT	Diphtheria toxin neutralization	404 (22-2816)	1,226 (352-11,264)
LT+DT	Anti-DT	4976 (669-46,909)	ND
CT+HIV p55 gag	Anti-p55	10,630 (1063-52,597)	ND
CT+Killed Rabies Virus	Anti-G protein	1.03 (IU/ml) (0.31-2.77)	7.54 (IU/ml) (3.31-17.47)

ND=not done. ELISA units (EU) shown as geometric mean and range in brackets.

## Example 21

### Langerhans cell activation

In two subjects, the site of immunization and the contralateral unimmunized arm were biopsied, one at 24 hours post-immunization and one at 48 hours after the second immunization. Hematoxylin and eosin (H&E) staining of the specimens confirmed the clinical findings suggesting that no inflammation was seen after the immunization (Fig 1 A,B). Although routine histologic sections were unremarkable, LCs visualized using anti-CD1a staining of specimens from the site of immunization demonstrated greatly enlarged cell bodies but otherwise normal numbers of cells when compared to the control biopsies from the opposite arm, both at 24 and 48 hours (Fig. 1 C,D,E,F). Similar findings were seen using anti-HLA-DR and anti-S-100 to visualize LCs (not shown). LC morphology in the TCI immunized skin was similar in appearance to tonsillar crypt LCs that are thought to be chronically activated by lipopolysaccharides from the flora of the mouth (Noble).

Because of the limited size and number of human skin biopsy specimens examined, complementary murine studies were performed. LC activation in murine systems using contact sensitizers, LPS, and proinflammatory cytokines is characterized by both changes in morphology (Aiba) and through elevations in surface marker expression (Jakob and Udey). Mouse ear skin is frequently used for studies of LC activation and has also been shown to be an excellent site for transcutaneous immunization (Scharton-Kersten). Epidermal sheets were prepared 24 hours after application of CT to the ear and stained for MHC class II, an LC-restricted marker in murine skin. In comparison to the PBS treated ears, Fig. 2A, LC in CT treated ears exhibited marked changes in LC morphology with loss of dendritic processes, enlarged cell bodies, and intense staining of the cells - features of LC activation (Aiba) (Fig. 2 B,C). The LC-activating potential of CT was confirmed using flow cytometry. LC from CT-treated skin expressed increased levels of MHC Class II antigens and CD 86 (B7-2) and decreased levels of E-cadherin, consistent with LC activation described elsewhere (Pierre, Aiba, Jakob).

Immunization Procedure which may be used for Example 22.

5 BALB/c mice may be shaved with a #40 clipper. This shaving could be done without any signs of trauma to the skin. The shaving could be done from the mid-thorax to just below the nape of the neck. The mice can then be allowed to rest for 24 hours. Prior to this the mice could be ear-tagged for identification, and pre-bled to obtain a sample of pre-immune serum. Mice could also be transcutaneously  
10 immunized without shaving by applying 5-500  $\mu$ l of immunizing solution to each ear. The mice could be immunized in the following way. Mice could be anesthetized with 0.03-0.06 ml of a 20 mg/ml solution of xylazine and 0.5 ml of 100 mg/ml ketamine and immobilized by this dose of anesthesia for approximately 1-3 hours. The mice could be placed ventral side down on a warming blanket.

15 The immunizing solution and penetration enhancement compound (or technique) could be placed on the dorsal shaved skin of a mouse in the following manner: a 1.2 cm x 1.6 cm stencil made of polystyrene is laid gently on the back and a saline-wetted sterile gauze could be used to partially wet the skin (allowing even application of the immunizing solution), the immunizing solution could then be  
20 applied with a pipet to the area circumscribed by the stencil to yield a 2 cm<sup>2</sup> patch of immunizing solution. Care could be used not to scrape or rub the skin with the pipet tip. The immunizing solution could be spread around the area to be covered with the smooth side of the pipet tip. Alternatively, the immunizing solution could be placed directly on the skin without wetting or with wetting without the use of a stencil.

25 The immunizing solution (between about 5  $\mu$ l and about 200  $\mu$ l) could be left on the back of the mouse for 60-120 minutes. At the end of the immunization period, the mouse could be held gently by the nape of the neck and the tail under a copious stream of lukewarm tap water, and washed for 10 seconds. The mouse could then be gently patted dry with a piece of sterile gauze and a second washing could then be  
30 performed for 10 seconds; the mouse could then be patted dry a second time and left in the cage. The mice would appear to exhibit no adverse effects from the anesthesia, immunization, washing procedure, or toxicity from the exotoxins. No skin irritation, swelling or redness would be seen after the immunization and the mice would appear

to thrive. Immunization using the ear could be performed as described above except that fur would not be removed prior to immunization.

#### Antigen

- 5           The following antigens could be used for immunization and ELISA, and could be mixed using sterile PBS or normal saline. Cholera toxin or CT (List Biologicals, Cat #101B, lot #10149CB), CT B subunit (List Biologicals, Cat #BT01, lot #CVXG-14E), CT A subunit (List Biologicals, Cat #102A, lot #CVXA-17B), CT A subunit (Calbiochem, Cat #608562); pertussis toxin, salt-free (List Biologicals, lot #181120a);
- 10   tetanus toxoid (List Biologicals, lots #1913a and #1915a); Pseudomonas exotoxin A (List Biologicals, lot #ETA25a); diphtheria toxoid (List Biologicals, lot #15151); heat-labile enterotoxin from E. coli (Sigma, lot #9640625); bovine serum albumin or BSA (Sigma, Cat #3A-4503, lot #31F-0116); and Hemophilus influenza B conjugate (Connaught, lot#6J81401).

15   ELISA - IgG(H+L)

Antibodies specific for CT, LT, ETA, pertussis toxin, diphtheria toxoid, tetanus toxoid, Hemophilus influenza B conjugate, and BSA could be determined using ELISA in a technique similar to Glenn et al. (1995).

## Example 22

Activation of LT may be performed by incubating the LT with trypsin (or  
5 trypsin immobilized on beads) with or without reducing agents (e.g., dithiotheritol) to  
break the disulphide bonds near the trypsin cleavage site, under standard reaction  
conditions. Native LT can be activated by incubation of 100 µg of protein with 0.1  
µg trypsin in a total reaction volume of 100 µl for 45 min at 37°C. Alternatively, the  
trypsin can be fixed to beads and the LT may be eluted over the trypsin beads.

- 10 Trypsin cleavage can be demonstrated by SDS-PAGE (Laemilli, U.K., 1970, Cleavage  
of structural proteins during the assembly of the head of bacteriophage T4, Nature  
227:680-685). LT either treated or not treated with trypsin can be mixed with buffer  
containing dithiothreitol, and heated to 100°C for 5 min prior to SDS-PAGE analysis.  
Trypsin-treated LT could have a proteolytic fragment of 21K daltons consistent with  
15 trypsin cleavage of the A1 and A2, allowing the A1 subunit to ADP-ribosylate G  
proteins and therefore exert its toxic effects whereas untreated LT would demonstrate  
a band at 28K daltons, consistent with an intact A subunit. Activation can further be  
demonstrated in the mouse Y-1 cell assay in which native LT would be 1,000 fold less  
active than CT, but trypsin-treated LT would be equally as active as CT. Activation  
20 can also be demonstrated using an enzymatic assay, the NAD:agmatine ADP-  
ribosyltransferase assay. In such an assay, non-trypsin-treated LT would be expected  
to show low or undetectable activity whereas trypsin-treated LT would be expected to  
show similar activity as that demonstrated by CT.

### Example 23.

Transcutaneous immunization may be more useful if the immunization can be performed over a short period of time. It may be useful for example for an immunization to be performed during a routine clinic visit lasting 30 minutes. In this example we show that transcutaneous immunization can be performed in hydrated, alcohol swabbed skin in such a short period.

C57BL/6 mice 6 to 8 weeks of age were anesthetized and shaved as described in the "immunization procedure". On the day of immunization the backs of the mice were wiped with isopropanol. After the alcohol had evaporated (approximately 10 minutes), 200  $\mu$ l of water was applied to the back for hydration. 15 minutes later the immunization solution was applied to the back and left for the specified period of time. Removal of excess antigen was conducted as described in the "immunization procedure." Mice were immunized with CT alone (100  $\mu$ g in 50  $\mu$ l) at d0 and with CT plus DT (100  $\mu$ g each in 100  $\mu$ l volume) at 4, 6 and 9 weeks. Twelve weeks after the primary immunization the animals were bled and the anti-DT titers determined using an ELISA as described above for "ELISA IgG (H+L)". The results are shown in Table 23.

Anti-DT titers were clearly elevated in serum from all of the animals immunized with CT and DT when compared with titers in serum from the same animals prior to immunization (prebleed). Maximal effects of immunization appeared to occur in animals vaccinated for a period of 60 minutes although the titers were similar at 30 and 120 minutes. Fifteen minutes of immunization seemed less efficient as the titers in this group were approximately 10 fold less than that observed in the 30, 60 and 120 minute groups. Thus it appears that TCI can be achieved within 15 minutes of antigen application.

Table 23. Effect of the duration of antigen application on humoral immunity induced by transcutaneous immunization.

ear tag #	duration of immunization	anti-DT IgG (ELISA units)		
		prebleed	12 weeks	geomean
361	15 min	6	214	300
362	15 min		664	
363	15 min		314	
364	15 min		181	
365	15 min		1594	
366	30 min	8	11953	13445
367	30 min		32478	
368	30 min		24346	
369	30 min		3457	
370	30 min		99776	
371	60 min	12	75787	107963
372	60 min		200768	
373	60 min		102592	
374	60 min		87034	
375	60 min		9210	
376	120 min	4	48132	48202
377	120 min		99362	
378	120 min		37308	
379	120 min		30255	
380	120 min		25149	

All publications referred to in this application are incorporated by reference herein as indicative of the state of the art.

## REFERENCES

5

Alving, C.R., and Wassef, N.M. (1994) Cytotoxic T lymphocytes induced by liposomal antigens: Mechanisms of immunological presentation. *AIDS Res. Hum. Retro.*, 10(sup. 2):S91-S94.

10

Antel, J.P., et al. (1996) Immunotherapy for multiple sclerosis: From theory to practice. *Nature Medicine*, 2:1074-1075.

Ausubel, F.M., et al. (1996) *Current Protocols in Molecular Biology*, Wiley, New York.

15

Bathurst, I.C., et al. (1993) An experimental vaccine cocktail for *Plasmodium falciparum* malaria. *Vaccine*, 11:449-456.

20

Blum, H.E. (1995) Variants of hepatitis B, C and D viruses: Molecular biology and clinical significance. *Digestion*, 56:85-95.

Bodanszky, M. (1993) *Peptide Chemistry*, Springer-Verlag, New York.

25

Bos, J.D. (1997) The skin as an organ of immunity. *Clin. Exp. Immunol.*, 107 (suppl. 1):3-5.

Burnette, W.N., et al. (1994) Recombinant microbial ADP-ribosylating toxins of *Bordetella pertussis*, *Vibrio cholerae*, and enterotoxigenic *Escherichia coli*: Structure, function, and toxoid vaccine Development. In: *Bioprocess Technology*, (Eds.

30

Burnette, W.N., et al.), pp. 185-203.



Chang, S.P., et al. (1989) Generalized immunological recognition of the major merozoite surface antigen (gp 195) of *Plasmodium falciparum*. *Proc. Natl. Acad. Sci. USA*, 86:6343-6347.

- 5 Chang, S.P., et al. (1992) A carboxyl-terminal fragment of *Plasmodium falciparum* gp195 expressed by a recombinant baculovirus induces antibodies that completely inhibit parasite growth. *J. Immunol.*, 139:548-555.

- Chang, S.P., et al. (1994) Regulation of antibody specificity to *Plasmodium falciparum* merozoite surface protein-1 by adjuvant and MHC haplotype. *J. Immunol.*, 152:3483-3490.
- 10

- Clements, J.D., and Finkelstein, R.A. (1979) Isolation and characterization of homogenous heat-labile enterotoxins with high specific activity from *Escherichia coli* cultures. *Infect. Immunol.*, 24:760-769.
- 15

- Craig, J. (1965) The effect of cholera stool and culture filtrates on the skin guinea pigs and rabbits, In: *Proceedings of the Cholera Research Symposium, Honolulu, US Public Health Service Publication No 1328*, pp. 153-158.
- 20

- Craig, J. (1972) Cutaneous Responses to Cholera Skin Toxin in Man. I. Responses in Unimmunized American Males, In: *The Journal of Infectious Diseases*, Vol. 125, No. 3, pp. 203-215.

- 25 Dahl, M.V. (1996) Atopic dermatitis. In: *Clinical Immunodermatology*, 3rd Ed. Mosby, St. Louis, pp. 345-352.

- Delenda, C., et al. (1994) Analysis of C-terminally truncated dengue 2 and dengue 3 virus envelope glycoproteins: Processing in insect cells and immunogenic properties in mice. *J. Gen. Virol.*, 75:1569-1578.
- 30

Deprez, B., et al. (1996) Comparative efficiencies of simple lipopeptide constructs for in vivo induction of virus-specific CTL. *Vaccine*, 14:375-382.

Deutscher, M.P. (1990) Guide to Protein Purification, Academic Press, San Diego.

5 Dickenson, B.L., and Clements, J.D. (1995) Dissociation of Escherichia coli heat-labile enterotoxin adjuvanticity from ADP-ribosyltransferase activity. Infect. Immun., 63:1617-1623.

Dragunsky, E.M., et al. (1992) Experimental evaluation of antitoxic protective effect of new cholera vaccines in mice. Vaccine, 10:735-736.

10

Elson, C.O., and Dertzbaugh, M.T. (1994) Mucosal adjuvants. In: Handbook of Mucosal Immunology (Eds. Ogra, P.L., et al.) Academic Press, San Diego, p. 391.

15

Finkelstein, R.A., and LoSpallutto, J.J. (1969) Pathogenesis of experimental cholera: Preparation and isolation of cholera toxin and cholera toxinoid, J. Exp. Med. 130:185-202.

Fonseca, B.A., et al. (1994) Recombinant vaccinia viruses co-expressing dengue-1 glycoproteins prM and E induce neutralizing antibodies in mice. Vaccine, 12:279-285.

20

Frankenburg, S., et al. (1996) Effective immunization of mice against cutaneous leishmaniasis using an intrinsically adjuvanted synthetic lipopeptide vaccine. Vaccine, 14:923-929.

25

Fries, L.F., et al. (1992a) Liposomal malaria vaccine in humans: A safe and potent adjuvant strategy. Proc. Natl. Acad. Sci. USA, 89:358-362.

30

Fries, L.F., et al. (1992b) Safety, immunogenicity, and efficacy of a Plasmodium falciparum vaccine comprising a circumsporozoite protein repeat region peptide conjugated to Pseudomonas aeruginosa toxin A. Infect. Immun., 60:1834-1839.

Glenn, G.M., et al. (1995) Murine IgG subclass antibodies to antigens incorporated in liposomes containing lipid A. Immunol. Lett., 47:73-78.

Glenn, G. M., Scharton-Kersten, T., Vassell, R., Mallet, C.P., Hale, T.L. & Alving, C.R. Transcutaneous immunization with cholera toxin protects mice against lethal mucosal toxin challenge. *J. Immunol.* **161**, 3211-3214 (1998).

- 5 Glenn, G.M., Rao, M., Matyas G.R. & Alving, C.R. Skin immunization made possible by cholera toxin. *Nature* **391**, 851 (1998).

Glenn, G.M., Scharton-Kersten, T., Vassell, R., Matyas, G. & Alving, C.R.  
Transcutaneous immunization using bacterial ADP-ribosylating exotoxins as antigens  
10 and adjuvants. *Infect. Immun.* (in the press).

Goeddel, D.V. (1990) Gene Expression Technology, Academic Press, San Diego.

- 15 Gordon, Ada and Alistar Ramsey (1997) Vaccine, Vaccination and the Immune Response, Philadelphia, New York: Lipincott-Raven, pp. 13-14.

Gregoriadis, G. (1993) Liposome Preparation and Related Techniques, 2nd Ed., CRC Press, Boca Raton.

- 20 Herrington, D.A., et al. (1991) Safety and immunogenicity of a recombinant sporozoite malaria vaccine against Plasmodium vivax. *Am. J. Trop. Med. Hyg.*, 45:695-701.

- 25 Hollingsbee, D. (1995) Use of hydrocolloid patches, In: Percutaneous Penetration Enhancers (Eds., Smith, E. and Maibach, H.), CRC Press.

Idson, B. (1978) Hydration and percutaneous absorption. *Curr. Prob. Dermatol.*, 7:132-141.

- 30 Jahrling, P.B., et al. (1996) Passive immunization of Ebola virus-infected cynomolgus monkeys with immunoglobulin from hyperimmune horses. *Arch. Virol. Suppl.*, 11:135-140.

Janeway, C.A., and Travers, P. (1996). Immunobiology, Churchill Livingstone, New York.

Janson, J.-C., and Ryden, L. (1989) Protein Purification, VCH, New York.

5

Katkov, W.N. (1996) Hepatitis vaccines. Med. Clin. North Am., 80:189-200.

Khusmith, S., et al. (1991) Protection against malaria by vaccination with sporozoite surface protein 2 plus CS protein. Science, 252:715-718.

10

Kounnas, M.Z., et al. (1992) The  $\alpha$ 2-macroglobulin receptor/low density lipoprotein receptor-related protein binds and internalizes Pseudomonas exotoxin A. J. Biol. Chem., 267:12420-12423.

15

Kriegler, M. (1990) Gene Transfer and Expression, Stockton Press, New York.

Krueger, K.M., and Barbieri, J.T. (1995) The family of bacterial ADP-ribosylating exotoxins. Clin. Microbiol. Rev., 8:34-47.

20

Lee, A., and Chen, M. (1994) Successful immunization against gastric infection with Helicobacter species: Use of a cholera toxin B-subunit-whole-cell vaccine. Infect. Immun., 62:3594-3597.

25

Leung, D.Y. (1997) Atopic dermatitis: Immunobiology and treatment with immune modulators. Clin. Exp. Immunol., 107 (Suppl. 1):25-30.

Lieberman, J.M., and Greenberg, D.P. (1996) Hepatitis A and B vaccines in children. Adv. Pediatr. Infect. Dis., 11:333-363.

30

Malik, A., et al. (1991) Human cytotoxic T lymphocytes against the Plasmodium falciparum circumsporozoite protein. Proc. Natl. Acad. Sci. USA, 88:3300-3304.

Mast, E.E., and Krawczynski, K. (1996) Hepatitis E: An overview. *Annu. Rev. Med.*, 47:257-266.

5 Mckenzie, A.W., and Stoughton, R.B. (1962) Method for comparing percutaneous absorption of corticosteroids. *Arch. Dermatol.*, 86:608-610.

Medzhitov, R., and Janeway, C.A. (1997) Innate immunity: Impact on the adaptive immune response. *Curr. Opin. Immunol.*, 9:4-9.

10 Mekalanos, J.J., et al. (1979) Enzymatic activity of cholera toxin II. Relationships to proteolytic processing, disulphide bond reduction, and subunit composition, *J. Biol. Chem.*, 254:5855-5861.

15 Migliorini, P., et al. (1993) Malaria vaccine: Immunization of mice with a synthetic T cell helper epitope alone leads to protective immunity. *Eur. J. Immunol.*, 23:582-585.

Morein, B., and Simons, K. (1985) Subunit vaccines against enveloped viruses: Virosomes, micelles and other protein complexes. *Vaccine*, 3:83-93.

20 Munoz, E., et al. (1990) Cholera toxin discriminates between T helper 1 and 2 cells in receptor-mediated activation: Role of cAMP in T cell proliferation. *J. Exp. Med.*, 172:95-103.

25 Murray, E.J. (1991) *Gene Transfer and Expression Protocols*. Humana Press, Clifton, New Jersey.

Nohria, A., and Rubin, R.H. (1994) Cytokines as potential vaccine adjuvants. *Biotherapy*, 7:261-269.

30 Paul, A., and Cevc, G. (1995) Noninvasive administration of protein antigens: Transdermal immunization with bovine serum albumin in transfersomes. *Vaccine Res.*, 3:145-164.

Paul, A., et al. (1995) Transdermal immunization with large proteins by means of ultradeformable drug carriers. *Eur. J. Immunol.*, 25:3521-3524, 1995.

5 Paul, W.E., and Seder, R.A. (1994) Lymphocyte responses and cytokines. *Cell*, 76:241-251.

Pessi, A., et al. (1991) Lack of H-2 restriction of the *Plasmodium falciparum* (NANP) sequence as multiple antigen peptide, *Eur. J. Immunol.*, 24:2273-2276.

10 Pierce, N.F. (1978) The role of antigen form and function in the primary and secondary intestinal immune responses to cholera toxin and toxoid in rats. *J. Exp. Med.*, 148:195-206.

15 Pierce, N.F., and Reynolds, H.Y. (1974) Immunity to experimental cholera. I. Protective effect of humoral IgG antitoxin demonstrated by passive immunization. *J. Immunol.*, 113:1017-1023.

Plotkin, S.A., and Mortimer Jr., E.A. (1994) *Vaccines*, 2nd Ed., W.B. Saunders, Philadelphia.

20 Proksch, E., and Brasch, J. (1996) Integrity of the permeability barrier regulates epidermal Langerhans cell density. *Br. J. Dermatol.*, 134:630-638.

25 Proksch, E., and Brasch, J. (1997) Influence of epidermal permeability barrier disruption and Langerhans' cell density on allergic contact dermatitis. *Acta Derm. Venereol.*, 77:102-104.

Rappuoli, R., et al. (1995) Genetic detoxification of bacterial toxins: A new approach to vaccine development. *Int. Archiv. Allergy Immunol.*, 108:327-333.

30 Rappuoli, R., et al. (1996) New vaccines against bacterial toxins. *Adv. Exp. Med. Biol.*, 397:55-60.

Ribi, H.O., et al. (1988) Three-dimensional structure of cholera toxin penetrating a lipid membrane. *Science*, 239:1272-1276.

5 Richards, R.L., et al. (1995) A compendium of vaccine adjuvants and excipients. In: *Vaccine Design* (Eds., Powell, M.F., and Newman, M.J.), Plenum, New York.

Roberts, M.S., and Walker, M. (1993) Water, the most natural penetration enhancer. In: *Pharmaceutical Skin Penetration Enhancement* (Eds., Walters, K.A., and Hadgraft, J.), Marcel Dekker, New York.

10

Saloga, J., et al. (1996) Superantigens. *Exp. Dermatol.*, 5:65-71.

Saukkonen, K., et al. (1992) Pertussis toxin has eukaryotic-like carbohydrate recognition domains. *Proc. Natl. Acad. Sci. USA*, 89:118-122.

15

Schneerson, R.E., et al. (1996) A toxoid vaccine for pertussis as well as diphtheria? Lessons to be relearned. *Lancet* 348:1289-1292.

Scopes, R.K. (1993) *Protein Purification*, Springer-Verlag, New York.

20

Seder, R.A., and Paul, W.E. (1994) Acquisition of lymphokine-producing phenotype by CD4<sup>+</sup> T cells. *Annu. Rev. Immunol.*, 12:635-673.

Shafara, A., et al. (1995) Hepatitis C. *Ann. Intern. Med.*, 125:658-668.

25

Skeiky, Y.A.W., et al. (1995) A recombinant *Leishmania* antigen that stimulates human peripheral blood mononuclear cells to express a Th1-type cytokine profile and to produce interleukin 12. *J. Exp. Med.*, 181:1527-1537.

30 Smedile, A., et al. (1994) Advances in hepatitis D virus biology and disease. *Prog. Liver Dis.*, 12:157-175.

Smucny, J.J., et al. (1995) Murine immunoglobulin G subclass responses following immunization with live dengue virus or a recombinant dengue envelope protein. *Am. J. Trop. Med. Hyg.*, 53:432-437.

- 5 Sniderman, D.P. (1995) The mucosal adjuvant activities of ADP-ribosylating bacterial enterotoxins. *Crit. Rev. Immunol.*, 15:317-348.

Spangler, B.D. (1992) Structure and function of cholera toxin and the related *Escherichia coli* heat-labile enterotoxin. *Microbiol. Rev.*, 56:622-647.

10

Stacey, K.J., et al. (1996) Macrophages ingest and are activated by bacterial DNA. *J. Immunol.*, 157:2116-2122.

15

Stingl, G., et al. (1989) The immune functions of epidermal cells. *Immunol. Ser.*, 46:3-42.

20

Streilein, J.W., and Grammer, S.F. (1989) In vitro evidence that Langerhans cells can adopt two functionally distinct forms capable of antigen presentation to T lymphocytes. *J. Immunol.*, 143:3925-3933.

Summers, M.D., and Smith, G.E. (1987) A manual of methods for baculovirus vectors and insect cell culture procedure. Texas Agricultural Experiment Station Bulletin, No. 1555.

- 25 Tam, J.P. (1988) Synthetic peptide vaccine design: Synthesis and properties of a high-density multiple antigenic peptide system. *Proc. Natl. Acad. Sci. USA*, 85:5409-5413.

Tang, D.C., et al. (1997) Vaccination onto bare skin. *Nature*, 388:729-730.

- 30 Tew, J.G., et al. (1997) Follicular dendritic cells and presentation of antigen and costimulatory signals to B cells. *Immunol. Rev.*, 156:39-52.



Trach, D.D., et al. (1997) Field trial of a locally produced, killed, oral cholera vaccine in Vietnam. *Lancet*, 349:231-235.

Udey, M.C. (1997) Cadherins and Langerhans cell immunobiology. *Clin. Exp. Immunol.*, 107 (Suppl. 1):6-8.

van Heyningen, W.E., and Seal, J.R. (1983) *Cholera: The American Scientific Experience, 1947-1980*, Westview Press, Boulder, Colorado.

10 Vandembark, A.A., et al. (1996) Treatment of multiple sclerosis with T-cell receptor peptides: Results of a double-blind pilot trial. *Nature Medicine*, 2:1109-1115.

15 Vreden, S.G.S., et al. (1991) Phase I clinical trial of a recombinant malaria vaccine consisting of the circumsporozoite repeat region of *Plasmodium falciparum* coupled to hepatitis B surface antigen, *Am. J. Trop. Med. Hyg.*, 45:533-538.

Wang, R., et al. (1995) Induction of protective polyclonal antibodies by immunization with a *Plasmodium yoelii* circumsporozoite protein multiple antigen peptide vaccine. *J. Immunol.*, 154:2784-2793.

20 White, K., et al. (1993) Induction of cytolytic and antibody responses using *Plasmodium falciparum* repeatless circumsporozoite protein encapsulated in liposomes. *Vaccine*, 11:1341-1346.

25 Wiesmueller, K.-H., et al. (1991) The antibody response in BALB/c mice to the *Plasmodium falciparum* circumsporozoite repetitive epitope covalently coupled to synthetic lipopeptide adjuvant. *Immunology*, 72:109-113.

Wisdom, G.B. (1994) *Peptide Antigens*, IRL Press, Oxford.

30 Zhang, T., et al. (1995) Oral immunization with the dodecapeptide repeat of the serine-rich *Entamoeba histolytica* protein (SREHP) fused to the cholera toxin B

[illegible]